

S/N 0102- LF- 014- 6601

407 997

2

SECURITY CLASSIFICATION OF THIS PAGE (When Date Ent

4 10


1	
-	
•	the second of th
✓	
• • • • • • • • • • • • • • • • • • • •	
	[2018] [2018] [2018] [2018] [2018] [2018] [2018] [2018] [2018] [2018] [2018] [2018] [2018] [2018] [2018] [2018]

DISCLAIMER NOTICE

THIS DOCUMENT IS BEST QUALITY PRACTICABLE. THE COPY FURNISHED TO DDC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

ABSTRACT

During our synthesis of phthalocyanatoruthenium(II) and phthalocyanatoosmium(II) complexes reported by Berizin and Sennikova, and by Krueger and Kenny we observed a remarkable solubility of these complexes in common organic solvents. The solubility of these complexes enabled us to isolate several pure reported and new phthalocyanine complexes. Among the complexes we studied carbonylphthalocyanato(pyridine)ruthenium(II)[PcRu(CO)(Py), carbonylphthalocyanatopyridineosmium(II)[PcOs(CO)(Py)], and carbonylphthalocyanato (tetrahydrofuran)ruthenium(II)[PcRu(CO)(THF)] were isolated in pure form. They are among the first reported metallophthalocyanines with a carbonyl as one of their axial ligands. Furthermore, several new ways of synthesizing these complexes in quantitative yields have been established. The structure of PcOs(CO)(Py) elucidated by the X-ray diffraction analysis. The osmium ion is octahedrally coordinated with the carbonyl and pyridine groups axially coordinated. The pyridine ring is tilted slightly with respect to the perpendicular to the phthalocyanine ring. The interplanar angle is 98.6°. Interesting comparisons may be made between PcOs(CO)(Py) and related porphyrin complexes;



Phthalocyanine and metallophthalocyanines demonstrate significant electrical and photo properties, 1,2 e.g. semiconductivity, photoconductiviety, photochemical reactivity, luminescence, and fluorescence, which are relevant to the current world problem of conversion and production of energy. The number of publications per year on chemistry of phthalocyanines and metallophthalocyanines has tripled during the past decade. During 1977, nearly 700 papers and patents were published. In spite of the increasing interest in this area and the near completion of the synthesis of all metal ions having the normal classic configuration, basic phthalocyanine chemistry has lacked the advantage of any dynamic or extraordinary progress. Many metalloporphyrins have been made and studied extensively, and a large body of coordination chemistry of these complexes has been developed, while the coordination chemistry of metallophthalocyanines has been poorly developed. This contrast is mainly due to the significantly lower solubility of metallophthalocyanines and has led to the presumption that the development of coordination chemistry of phthalocyanines is difficult in general.

During our synthesis of pthalocyanatoruthenium(II) and phthalocyanatoosmium(II) complexes reported by Berizin and Sennikova^{4,5} and by Krueger and
Kenny,^{6a} we have found a remarkable solubility of these complexes in common
organic solvents. The solubility of these complexes enabled us to isolate
several pure reported and new phthalocyanine complexes. Among the complexes
we studied, carbonylphthalocyanato(pyridine)ruthenium(II)[PcRu(CO)(Py)],
carbonylphthalocyanato(pyridine)osmium(II)[PcOs(CO)(Py)], and carbonylphthalocyanato(tetrahydrofuran)ruthenium(II)[PcRu(CO)(THF)] were isolated in pure form.
These compounds are among the first reported metallophthalocyanines with carbonyl
as one of their axial ligands. Furthermore, several new ways of synthesizing these

complexes in quantitative yields have been established. Since the completion of this work, ^{6c} the preparation of PcRu(CO)(L)L (L=Py, 4-MePy; 4-t-BuPy) has been reported by N. P. Farrell et al. ^{6b} However, the methods of preparation described herein, which differ from their techniques, produce products in considerably higher yield. In this paper we also describe the structure of PcOs(CO)(Py)Py as determined from an x-ray diffraction analysis. This is the first reported structure of this type of phtalocyanine complex and provides interesting comparisons with analogous porphyrin complexes.

Experimental Section

Materials

Phthalonitrile (practical) was purchased from the Eastman Kodak Co. and used without any further purifications. Ruthenium-trichloride-(trihydrate)(RuCl₃.3(H₂O)) and osmium tetraoxide(OsO₄) were purchased from the Ventron Corporation and used without any further purifications. Ruthenium dodecacarbonyl(Ru₃(CO)₁₂) and osumium dodecrabonyl(Os₃(CO)₁₂) were purchased from Strem Chemicals and used without purification. Carbon monoxide gas (99.9%) was purchased from the Matheson Gas Company and used without purification. Pyridine (reagent grade), tetrahydrofuran (reagent grade), methylene chloride (spectro grade), chloroform (spectro grade), and benzene (spectro grade) were used without further purification. Neutral alumina (60 - 100 mesh), acid alumina (80 - 200 mesh) and silica gel (60 - 200 mesh) were purchased from the Fisher Scientific Company.

Physical Measurements

Elemental analysis were performed by the Schwarzkopf Microanalytical Laboratory, N.Y. Visible spectra were measured with a Beckman Spectrophotometer Model 24. Infrared spectra were measured with a Beckman Infrared Spectrophotometer Model IR-8.

Isolation of PcRu(CO)(THF) from the Products of the Reaction of RuCl₃·3(H₂O) with Phthalonitrile

RuCl₂·3H₂O (1.00 g; 3.8 mmol) was heated with excess phthalonitrile (8.00 g; 62.5 mmol) at 250°C for 4 hours. After washing the resulting product with methanol several times, unreacted phthalonitrile was removed by sublimation at 150°C under vacuum. The blue-black residue (4.1 g), which was left at the bottom of the sublimator, was dissolved in aniline (10 mL), and metal-free phthalocyanine was removed from the solution by filtration. The aniline solution was poured into benzene (1.0 L) with stirring. A black precipitate (0.8 g) was separated by filtration from the benzene solution. The solution was then concentrated and dried under vacuum at 100°C until aniline in the solution was removed. The dried residue (3.5 g) was dissolved in benzene (30 mL), and then the solution was chromatographed over silica gel with benzene. Meanwhile, benzene was replaced by chloroform, and a light blue band, which was most intense, was separated from the dark blue band. The light blue eluate was condensed and chromatographed over neutral alumina with chloroform. Another bright blue band followed the original blue band when tetrahydrofuran was added to chloroform. The second blue eluate was condensed and dried. Fine needle crystals (120 mg) with a typical phthalocyanine appearance of red reflection and blue transmission were obtained. The visible spectrum of this compound shows maxima (in nm) at 642 and 581 (in chloroform). The infrared spectrum has an intense peak at 1960 cm -1.

Preparation of PcRu(CO)(Py) from RuCl₃·3(H₂O) and Phthalonitrile in Carbon Monoxide

RuCl₂·3(H₂O) (100 mg; 0.38 mmol) was heated in molten phthalonitrile (500 mg; 1.9 mmol) under carbon monoxide atmosphere at 250°C for 4 hours. After cooling the product to room temperature, pyridine (5 mL) was added to the cake. The mixture was again heated to the boiling temperature of pyridine for I hour. The pyridine solution of the resulting product was cooled to room temperature, and then pyridine was removed first by distillation. Excess amounts of phthalonitrile were removed by sublimation at 150° under vacuum. The blue residue (265 mg) was dissolved in methylene chloride (15 mL), and the solution was chromatographed over neutral alumina with methylene chloride. The first blue band was followed by a second blue band when methylene chloride was replaced with chloroform. The second blue eluate, which was present in large quantity, was condensed and dried at 100°C under vacuum. A fine blue powder (230 mg) with red reflection (PcRu(CO)(Py)) was obtained (84% yield). Anal. Found: Ru, 13.97; N, 17.35. Calc. for PcRu(CO)(Py): Ru, 14.02; N, 17.49. The visible spectrum shows maxima (in nm) at 624 and 581 (in chloroform). The infrared spectrum shows an intense peak at 1965 cm -1.

Preparation of PcRu(CO)(Py) from Ru₃(CO)₁₂ and Phthalonitrile

Ru₃(CO)₁₂ (100 mg; 0.16 mmol) was added to molten phthalonitrile (500 mg; 1.9 mmol) in air and heated at 250°C for 4 hours. The product was heated with pyridine (5 mL) at 150°C for 1 hour. Pyridine

and unreacted phthalonitrile were removed, and the residue was dissolved in methylene chloride. Chromatography of the solution over neutral alumina gave a single blue band with chloroform. The chloroform eluate was condensed and dried. The blue powder (PcRu(CO)(Py)) with red reflection weighed 315 mg (91% yield).

Isolation of PcOs(CO)(THF) from the Products of the Reaction of OsO₄ with Phthalonitrile

 OsO_{L} (1.0 g; 3.9 mmol) was added to molten phthalonitrile (6.0 g; 46.8 mmol) in a 50 mL flask equipped with a condenser. The mixture was kept at 250°C for 4 hours. The excess phthalonitrile was removed from the product by sublimation. The residue was put in an extraction thimble, and some products were extracted with tetrahydrofuran. The THF solution was condensed and dissolved in chloroform. The chloroform solution was chromatographed over acid alumina with chloroform. The first blue band was collected and condensed. The condensed residue was then chromatographed over neutral alumina with chloroform. The first blue band of the second chromatography was collected and condensed. After drying at 100°C under vacuum, a blue powder (PcOs(CO)(THF)) was obtained (115 mg; 3.7% yield). Anal. Found: Os, 23.20; N, 14.15. Calc. for PcOs(CO)(THF): Os, 23.69; N, 13.96. The visible spectrum of the compound shows maxima (in nm) at 636, and 576 in THF. The infrared spectrum of the compound shows an intense peak at 1930 cm⁻¹.

Preparation of PcOs(CO)(Py) from OsO, and Phthalonitrile in Carbon Monoxide

OsO, (107 mg; 0.42 mmol) was added to molten phthalonitrile (500 mg; 3.9 mmol) in a 50 mL round bottom flask equipped with a condenser under carbon monoxide stream. The mixture was kept at 250°C for 4 hours under a carbon monoxide atmosphere. The mixture was then cooled to room temperature and dissolved in pyridine (5 mL). The pyridine solution was refluxed for 1 hour under a carbon monoxide atmosphere. Pyridine and phthalonitrile were removed under vacuum from the product. The resulting blue residue was dissolved in chloroform (50 mL), and metal-free phthalocyanine was removed from the chloroform solution by filtration. The chloroform solution was condensed and chromatographed over neutral alumina with chloroform. The single blue band was collected and condensed. A blue powder with red reflection(PcOs(CO)(Py)) was obtained after drying (280 mg; 83% yield). Anal. Found: Os, 22.96; N, 15.48. Calc. for PcOs(CO)(Py): Os, 23.49; N, 15.57. The visible spectrum of the compound shows maxima (in nm) at 632 and 575 (in chloroform). The infrared spectrum of the compound shows an intense peak at 1930 cm -1.

Preparation of PcOs (CO) (Py) from Os (CO) 12 and Phthalonitrile

Os₃(CO)₁₂ (100 mg; 0.11 mmol) was added to molten phthalonitrile (500 mg; 3.9 mmol) in a 50 mL round bottom flask. The mixture was kept at 250°C for 4 hours. Pyridine (5 mL) was added to the mixture, and the pyridine solution was refluxed for 1 hour. Pyridine and excess

phthalonitrile were removed from the product by distillation and sublimation. The blue residue was then dissolved in chloroform, and metal-free phthalocyanine was separated from the solution by filtration. The chloroform solution was condensed and chromatographed over neutral alumina with chloroform. The single blue band was collected and condensed. After drying at 100°C under vacuum, a blue powder with red reflection (PcOs(CO)(Py)) was obtained (220 mg; 82% yield).

X-ray Study

Crystals of PcOs(CO)(Py) were grown from a chloroform solution. The crystal chosen for intensity measurements was a parallellopiped bounded by $\{100\}$, $\{010\}$ and $\{001\}$. The dimensions were $0.20 \times 0.20 \times 0.09$ mm in the direction of \underline{a} , \underline{b} and \underline{c} respectively. It was mounted in a capillary 7 at an arbitrary orientation, but with \underline{a} approximately parallel to the spindle axis.

Crystal Data are listed in Table 1. An Enraf-Nonius CAD-4 computer controlled diffractometer was used. The radiation (Mo K α , λ =0.71069A) was monochromatized by pyrolitic graphite. The instrument centered the crystal automatically. The setting angles for 25 reflections, measured at + and - 20, were used to index the cell and then were refined to give an orientation matrix, cell constants, and a Niggli matrix which indicated that the system was monoclinic. The systematic absences uniquely determined the space group.

The diffracted intensities were collected using the 0-20 scan technique. Scan speeds, which were determined by a rapid preliminary scan, ranged from 0.28 to 3.35 deg/min. depending on the intensity. Very weak reflections were measured at the maximum rate. The scan range for each reflection was equal to 0.90 + 0.35 tan 0. Other experimental conditions are described elsewhere. No evidence of crystal decomposition or machine instability was noted.

Independent reflections (4596) were measured out to a sin0 value of 0.54 or 22.5° in 0. Of these, 2820 had a net intensity greater than $2\sigma_{I}$ and were used in analysis. The standard deviation σ_{I} was defined in terms of the statistical variances of the counts as $\sigma_{I}^{2} = \sigma_{I(\text{count})}^{2} + (9.02I)^{2}$ $\sigma_{I(\text{count})}$ is the variance determined solely from counting statistics. Structure factors were calculated in the usual way, including correction for partial polarization of the incident beam due to the use of a monochromator.

Determination and Refinement of the Structure

Because there are four molecules in the unit cell of space group $P2_1/n$, all atoms lie in general positions. The position of the osmium atom was found from an unsharpened Patterson synthesis. The rest of the 49 nonhydrogen atoms were found from a series of difference syntheses. Least squares refinement using full matrix methods was carried out minimizing the function Σ w($|F_0| - |F_c|$)², where w = $1/\sigma_F^2$. Initially isotropic temperature factors were used, but in the final refinements all nonhydrogen atoms were varied assuming anisotropic thermal motion. The positions of 18 of the 21 hydrogen atoms could be found from ΔF maps. However, refinement of the hydrogen atom parameters led to chemically unreasonable bond lengths and angles. Hence the positions of all the hydrogen atoms were calculated (C-H = 0.95A) and their contributions included in the structure factor calculations, assuming an isotropic temperature factor, B, of 4.0A². The hydrogen atom parameters were not refined. The refinement converged with $R = \Sigma ||Fo| - |Fc||/\Sigma |Fo| = 0.031$ and $R_w = (\Sigma w(|F_0| - |F_c|)^2/\Sigma w|Fo|^2)^{\frac{1}{2}} = 0.035$.

A correction for anomalous dispersion was made for all nonhydrogen atoms. Scattering factors were from Ref. 12. The osmium atom was assumed to be in the zero ionization state. No evidence of secondary extinction was found.

Attempts to apply absorption corrections were made. Transmission coefficients, calculated using a Gaussian integration method (6x4x6 grid), varied from 0.43 to 0.68 with most being about 0.6. The R factors increased substantially (R=0.049, R =0.056) when refinements were attempted using the corrected data. No improvement was noted in the standard deviations. It was concluded that the errors introduced in applying the corrections to a relatively small crystal enclosed in a capillary were larger than those introduced by ignoring absorption effects. Thus, the final refinements were carried out on uncorrected data.

In the last cycle of refinement all shifts on positional and thermal parameters were less than one standard deviation, with the largest shift being 0.37 standard deviations. The final value of the standard deviation of an observation of unit weight, defined as $\left[\Sigma_W(|F_0|-|F_c|)^2/(N_0-N_V)\right]^{\frac{1}{2}}$ was 0.944 for $N_0=2820$ reflections and $N_V=442$ parameters.

There were two peaks of about 1.5 e/A³ in the final difference Fourier. These were quite close to the osmium atom. Neither they nor any of the other peaks were considered physically significant.

Most calculations were performed on a PDP 11/40 computer using the Enraf-Nonius structure determination package (SDP). Johnson's ORTEP, some molecular geometry calculations (using XANADU by Roberts and Sheldrick) and local programs were run on an Amdahl 470v/6 computer. Use was made of the PDP 11/40-Vector General graphics system. Data reduction was performed on the Honeywell computer at the University of Houston.

The final positional and thermal parameters are given in Table II. The final calculated positions of the hydrogen atoms are given in Table III.

Tables IV and V contain the root-mean-square components of thermal displacement along the principal axis of the thermal ellipsoids and the observed and calculated structure factors respectively. Tables III-V are available as supplementary material.

Results and Discussion

The following reactions for the synthesis of phthalocyanatoruthenium complexes have been carefully examined, and considerable discrepancies from the reported results have been found. Furthermore, a new carbonyl complex of ruthenium phthalocyanine has been isolated from the reaction products. The reaction of RuCl₃·3(H₂O) with phthalonitrile was reported to give PcRu(III)Cl as the major product of the reaction by Berizin and Sennikova, 4,5 as shown in the following equation:

$$RuCl_3(H_2O) + C_6H_4(CN)_2 + PcRu(III)C1$$

The reaction of $RuCl_3$ $^3(H_20)$ with o-cyanobenzoamide was reported to give PcRu(II) as its major product by Krueger and Kenny, ^{6a} as shown in the following equation:

$$RuCl_3 \cdot 3(H_2O) + NCC_6H_4CONH_2 \rightarrow PcRu(II)$$

From our investigations both of these reactions have been found to give PcRu(II) as the major product and PcRu(III)Cl as the minor one.

In addition, a small quantity of PcRu(II)(CO)L (L is a solvent used for the isolation) was detected in the crude PcRu(II) 6C₆H₅NH₂ according to its infrared spectrum, and was isolated by column chromatography. The yield of PcRu(II)(CO)L was less than 5%. Thus, the reactions above should be expressed by the following equations:

 $RuCl_3: 3(H_2O) + C_6H_4(CN)_2 \rightarrow PcRu(II) + PcRu(III)C1 + PcRu(II)(CO)L$ $RuCl_3: 3(H_2O) + NCC_6H_4CONH_2 \rightarrow PcRu(II) + PcRu(III)C1 + PcRu(II)(CO)L$ The yield of 5% for CO complex is rather surprising in two respects. In the first place, the source of carbon monoxide is unknown. Since the reaction of RuCl₃ 3(H₃O) with phthalonitrile was carried out in air without solvent there should not be any direct carbonyl sources in the system. In the second place, the major product was four-coordinate PcRu(II), whereas it appears that stable Ru(II) porphyrin complexes are generally six coordinate. 15-20

This carbonyl complex of ruthenium phthalocyanine can be synthesized in tow different ways directly in high yield (80-90%). The reaction of ${\rm RuCl}_3$ '3(H₂0) with excess phthalonitrile under carbon monoxide atmosphere gives an almost quantitative yield even after the isolation of the compound through column chromatography. The reaction of ${\rm Ru}_3({\rm CO})_{12}$ with excess phthalonitrile also gives a high yield of PcRu(II)(CO)(Py). This reactions are shown in the following equations:

$$RuC1_{3} \cdot 3(H_{2}O) + C_{6}H_{4}(CN)_{2} + CO \rightarrow PCRu(II)(CO)L \xrightarrow{pyridine}_{C^{D}u(II)(CO)(Py)}$$

$$Ru_{3}(CO)_{12} + C_{6}H_{4}(CN)_{2} \longrightarrow RcRu(II)(CO)L \xrightarrow{pyridine}_{PcRu(II)(CO)(Py)}$$

The carbonyl complex of ruthenium phthalocyanine was characterized by infrared and visible absorption spectroscopy and elemental analysis. The infrared spectrum of the compound shows an intense band at 1965 cm⁻¹ which is assigned to $vC\equiv 0$ attached to ruthenium ion.

A well-defined carbonyl complex of osmium phthalocyanine was also isolated from the products of the reported reactions of $0sO_4$ with molten phthalonitrile. Although the yield of PcOs(II)(CO)L is less than 5%, it it isolated as a pure complex by column chromatography. The isolation was made possible by the remarkable solubility of the complex in common organic solvents. The presence of a carbonyl ligand attached to osmium metal was evidenced by an intense band at 1930 cm⁻¹ in the infrared spectrum. This new type of compound, PcOs(II)(CO)L, coordinates another donor molecule as its last axial ligand. For example, tetrahydrofuran and pyridine can coordinate to the open sixth site of osmium to form a stable compound with octahedral configuration.

These carbonyl complexes of osmium phthalocyanine can be prepared in high yield (80-90%) by the reaction of OsO_4 with molten phthalonitrile in carbon monoxide atmosphere or the reaction of $Os_3(CO)_{12}$ with phthalonitrile in air. Subsequent treatment with pyridine or THF yield PcOs(II)(CO)(Py) and PcOs(II)(CO)(THF) respectively. Column chromatography gives a single blue band of each complex. An x-ray diffraction analysis has been carried out on the former complex.

Carbonyl complexes of ruthenium and osmium porphines have been synthesized and characterized by a number of workers. $^{15-20}$

Table VI summarized the carbonyl stretching frequencies of these complexes together with those of the carbonyl complexes of ruthenium and osmium phthalocyanocines.

(Tetraphenylporphinato)(carbonyl)(pyridine)uthenium(II) (TPPRu(CO)(Py)) shows a carbonyl peak at 1939 cm⁻¹, which is slightly lower than that (1965 cm -1P of PcRu(CO)(Py). Therefore, the back donation of 4d electrons of ruthenium to the anti-bonding π -orbitals of the carbonyl ligand seems to be less in the phthalocyanine complex than in the TPP complex. The same phenomenon is observed in osmium complexes. (Octaethylporphinato)(Carbonyl)(pyridine)Os(II) (OEPOs(CO)(Py)) shows a carbonyl peak at 1902 cm⁻¹, while PcOs(CO)(Py) shows a carbonyl peak at 1930 1. In this case back donation of 5d electrons from osmium to the anti-bonding π -orbital of the carbonyl ligand seems to be less in the phthalocyanine than in the OEP complex. Probably the structural difference of the rings causes the difference in the degree of the back donoation of d-electrons from the central metal ions to the coordinated carbon monoxide. Apparently, Ru(II) or Os(II) in the phthalocyanine carbonyl complexes donate less electron density to the carbonyl ligand than those in porphyrin rings.

It should also be noted that the PcRu(II) complex, which does not have a carbonyl ligand, is capable of coordinating carbon monoxide at room temperature under 1 atmosphere of carbon monoxide. PcRu(II) can be prepared by removing aniline molecules from PcRu(II) $\cdot 6(C_6H_5NH_2)$ under vacuum at high temperature (200°C) and it can form a carbonyl complex

in THF or pyridine by bubbling CO gas into the solution for one day. The original $PcRu(II) \cdot 6(C_6H_5NH_2)$ can be recovered by refluxing PcRu(II) in freshly distilled aniline for a few minutes as shown in the following equations:

PcRu(II)·6(C₆H₅NH₂)
$$\xrightarrow{\Delta$$
, vacuum

PcRu(II) CO in pyridine

PcRu(II) (CO) (Py)

CO in THF

PcRu(II) (CO) (THF)

The carbonylation process can be traced by noting changes in its visible spectrum as shown in Fig. I.

PcRu(II) is expected to demonstrate an interesting chemistry, because it does not possess a carbonyl ligand. On the other hand, the chemistry of ruthenium porphyrin is rather limited becasue removal of the carbonyl ligand is extremely difficult. ^{16,21} PcRu(II) appears to have the potential to form new ruthenium phthalocyanine complexes with various other molecules such as O₂, N₂, NO, and olefins. Also, the remarkable solubility of ruthenium and osmium phthalocyanine complexes will possibly open a new era in the coordination chemistry of metallophthalocyanines.

DESCRIPTION OF STRUCTURE OF PcOs(CO)(Py)

The structure of (carbonyl)(pyridine)phthalocyanatoosmium(II) is shown in Figure 2. Stereoviews are shown in Figures 3 and 4. The osmium ion is octahederally coordinated with the carbonyl and pyridine groups axially coordinated. The pyridine ring is tilted slightly with respect to the perpendicular to the phthalocyanine ring. The interplanar angle is 98.6°.

The structure of an analogous osmium porphyrin complex has not been reported, but the structure of (carbonyl)(pyridine)tetraphenylporphinatoruthenium(II), TPPRu(CO)(Py) has been published. Osmium and ruthenium have similar covalent radii. The phthalocyanine complex bears a strong resemblance to this metalloporphine. In addition, the structure of a carbonyl osmium porphodimethene complex with pyridine as the other axial ligand, OEPMe2Os(CO)(Py) has been reported. Interesting comparisons may be made between these two complexes and PcOs(CO)(Py).

Bond lengths and angles are given in Table VII. The average interatomic distances between the osmium atom and the isoindole nitrogen atoms of the phthalocyanine molecule is 2.01A. However, it should be noted that the bond lengths for Os-N(2) and Os-N(6) are 1.98A, while the other two distances are 2.03A. The difference corresponds to ~5 standard deviations, so it is statistically significant. The C-N-C and N-C-C angles involving the isoindole nitrogen atoms also differ in the two pairs of isoindole groups. Thus, for groups 1 and 3 (containing N(2) and N(6), the average C-N-C angle is 112°, while the N-C-C an is 107°. In groups 2 and 4 (containing N(4) and N(8)) the corresponding angles are 107° and 111°. The differences once again correspond to about 4 or 5 standard deviations and are therefore significant. We are unable to offer a plausible explanation for these differences, which are not observed in other phthalocyanine complexes.

The size of the central "hole" in phthalocyanine complexes has been estimated to be 0.046 -0.050A smaller than in corresponding porphyrin complexes. A similar difference is noted in the present case. The osmium atom is slightly out of the plane of the phthalocyanine molecule, so the size of the hole has to be estimated from the distances between opposite isoindole nitrogen atoms. These distances are 4.06 and 3.96A, corresponding to an average radius (Ct-N distance) of 2.01A for the macrocyclic hole. In TPPRu(CO)(Py)²² the corresponding distance is 2.05A. The M-N distances are correspondingly longer. In both TPPRu(CO)(Py)²² and OEPMe₂Os(CO)(Py)²⁴ these average 2.06A.

The Os-N bond lengths involving the isoindole nitrogen atoms are short for such bonds. On the other hand the Os-N_{Py} (N_{py} is the pyridine nitrogen atom) distance of 2.209(9)A is relatively long. Ru-N_{py} distances of 2.06-2.09Å have been reported. M-NH₃ bond lengths in amine complexes of Os(II) and Ru(II) range from 2.12-2.14A. He M-N_{py} distance in both TPPRu(CO)(Py) and OEPMe₂Os(CO)(Py) is also elongated (2.193(4) and 2.230(4)A respectively.)

This long bond may be due to steric interactions between atoms in the pyridine ring, particularly the \sim hydrogen atoms, and atoms in the macrocycle. This would block a closer approach to the osmium atom by the pyridine molecule. The osmium atom lies 0.15A on the other side of phthalocyanine molecule. However, the closest contacts involving the \sim hydrogen atoms are between H(33) and C(32) and between H(37) and C(16). These contacts are both about 2.73A, which is not an unusually short contact distance in such cases. A tabulation of such contacts in porphyrin complexes containing planar axial ligands shows a range of 2.45-2.9A. In TPPRu(CO)(Py) the closest contact is 2.51A.

The angle between the plane of the pyridine ring and the plane defined by N(2), Os, and N(6) is 48° . An angle of $^{\sim}45^{\circ}$ minimizes steric interaction.

whereas an angle of $~0^{\circ}$ would bring the ~hydrogen atoms into close contact with the isoindole nitrogen atoms. 31

It has also been suggested a lengthening of ~0.1A will occur in the axial M-N bond due to the trans effect of the carbonyl group. ²² In OEPRu(Py)₂ ³² the observed axial bond lenghts are indeed shorter, 2.09-2.10A, in spite of some relatively close contacts of the ~-hydrogen atoms of the pyridine ligands with atoms in the macrocycle. ³⁰ An even shorter M-N bond length of 2.00A has been reported in PcFe(Py)₂. ³³

The metal carbonyl distance of 1.83(1) is in good agreement with the values found in TPPRu(CO)(Py) OEPMe₂Os(CO)(Py). The Os-C-O angle is linear (177°).

Least squares plane of interest are given in Table VIII. As mentioned previously, the osmium atom is 0.15A out of the plane defined by the phthalocyanine group in a direction toward the carbonyl group. The displacement decreases to 0.099A if one considers only the plane of the four isoindole nitrogen atoms. By way of comparison, the ruthenium atom in TPPRu(CO)(Py) is 0.079A out of the plane of the porphyrin skeleton toward the carbonyl group.

In the discussion on TPPRu(CO)(Py) the authors attributed the out-of-plane displacement of the metal ion to either very strong metal-carbonyl bonds or to the inability of the large metal ion to fit into the plane of the macrocycle. The former possibility now appears more likely, since in OEPRu(Py)₂ the ruthenium(II) ion lies in the plane of the macrocycle. The same factor is probably primarily responsible for the out-of-plane displacement observed for the metal ion in this phtalocyanine complex.

The difference of 0.065A in the displacements of the metal ion from the planes of the macrocycle and of the isoindole nitrogen atoms indicate that the macrocycle itself is "domed", corresponding to a C_{4v} deviation from planarity. Deviations of equal or larger magnitude have been observed in other metallo-

phthalocyanine complexes, where the metal ion is out of the plane of the macrocycle. 34-37 The difference between the deviation of the osmium atom from the plane of the four isoindole nitrogen atoms and the plane defined by the pyrrole rings and the bridging nitrogen atoms is 0.027A, a value falling in the middle of the range found for porphyrins with the metal ion out of the macrocyclic plane. 38

The doming is not equal for the four isoindole groups. The maximum deviations from the plane of the four isoindole nitrogen atoms are 0.23 and 0.35A for phenyl carbon atoms in groups 2 and 3, while the maximum deviations in the same direction in groups 1 and 4 are 0.04 and 0.12A. Such a pattern was observed in aquophthalocyanatomagensium(II).

There appears to be some variation in bond parameters of phthalocyanine complexes as the size of the central "hole" increases. ³⁹ The C-N-C angle involving the azamethine nitrogen atom is the most sensitive bond parameter. This angle ranges from 121.7° in Fe(Pc)²⁵ (Ct-N: 1.93A) to 126.2° in Cl₂Sn(Pc)⁴⁰ (Ct-N: 2.05A). The average value of 125(1)° found for this angle in PcOs(CO)(Py) agrees with that found in phthalocyanine complexes with similar Ct-N distances. ^{35,39} The presence of a very heavy metal atom like osmium decreases the accuracy with which the lighter atoms can be determined, but within the observed standard deviations, the other bond parameters agree with those reported in the accurately determined structure of Zn(Pc)³⁹ and Sn(Pc). ³⁴

A packing diagram of the unit cell is shown in Figure 3. Table IX lists the intermolecular contacts $\leq 3.5A$. Most of the shortest contacts involve the carbonyl oxygen atom. Non-bonded contacts of this magnitude have been observed in other carbonyl complexes (e.g. $[H_3OEP]^+$ $[Re_2(CO)_6Cl_3]^-$). Weither these nor any of the other intermolecular contacts are believed to have any significant effect on the structure.

The structure of PcRu(CO)(Py) probably possesses a structure similar to that of OsPc(CO)Py.

Acknowledgment. This work was supported in part by the Office of Naval Research and the Robert A. Welch Foundation (Grant A-328) and the Texas Agricultural Experiment Station (Grant H-1668).

Supplementary Material Available: Table III-V containing calculated hydrogen atom positions, root-mean-square components of thermal ellipsoids and observed and calculated structure factors (pages). Ordering information is given on any current masthead page.

REFERENCES

- 1. Lever, A. B. P. "Advances in Inorganic Chemistry and Radiochemistry, Vol. 7, Academic Press: New York, 1965, p. 27.
- 2. Moser, F. H.; Thomas, A. L., "Phthalocyanine Compounds", ACS Monograph 157, Reinhold: New York, 1963.
- 3. Smith, K. M., Ed., "Porphyrins and Metalloporphyrins", Elsevier: Amsterdam, 1975.
- 4. Berizin, B. D.; Sennikova, G. V. Dokl. Akad. Nauk SSSR 1964, 159, 117.
- 5. Berizin, B. D.; Sennikova, G. V. Dokl. Akad. Nauk SSSR 1962, 146, 604.
- a. Krueger, P. C.; Kenny, M. E. <u>Inorg. Nucl. Chem.</u> 1963, 25, 303.
 b. Farrell, N. P.; Murray, A. J.; Thornback, J. R.; Dolphin, D. H.; James, B. R.; <u>Inorg. Chim. Acta.</u> 1978, 28 L 144. c. Omiya, S. Ph.D. Dissertation, <u>Texas A&M Univ.</u>, April, 1978.
- 7. Meyer, E. F., Jr. J. Appl. Crystallogr. 1973, 6, 45.
- 8. Niggli, P. "Handbuch der Experimentalphysik", Akademische Verlagsgesellschft, 1928; p. 108.
- 9. Instruction Manual, CAD-4 System, Enraf Nonius, Delft, 1977.
- Reisner, G. M.; Bernal, I.; Brunner, H.; Muschiol, M. <u>Inorg. Chem.</u> 1978, 17, 783.
- 11. Cromer, D. T.; Liberman, D. J. Chem. Phys. 1970, 53, 1891.
- 12. Cromer, D. T.; Waber, J. T. in "International Tables for X-Ray Crystallography", Vol. IV, Ibers, J. A.; Hamilton, W. C. Ed., Kynoch Press: Birmingham, England, 1974; pp 72-101.
- 13. Johnson, C. K. "ORTEP", Report ORNL-3794, Oak Ridge National Laboratory, Oak Ridge, Tenn., revised 1965.
- 14. Morimoto, C. N.; Meyer, E. F. Jr. in "Crystallographic Computing Techniques", Ahmed, F. R., Ed.; Munksgaard, Copenhagen, 1976.
- 15. Chow, B. C.; Cohen, I. A. Bioinorg. Chem. 1971, 1, 57.
- 16. Hopf, F. R.; Whitten, G. J. Am. Chem. Soc. 1976, 98, 7422.
- Srivastava, T. S.; Hoffman, L.; Tsutsui, M. J. Am. Chem. Soc. 1972, 94, 1385.
- Bonnet, J. J.; Eaton, S. S.; Eaton, G. R.; Holm, R. H.; Ibers, J. J. Am. Chem. Soc. 1973, 95, 2141.
- 19. Buchler, J. W.; Rohbock, K. J. Organometal. Chem. 1974, 65, 223.
- 20. Buchler, J. W.; Folz, M. Z. Naturforsch. 1977, 32b, 1439.

- 21. Farrell, N.; Dolphin, D. H.; James, B. R. J. Am. Chem. Soc. 1978, 100, 324.
- 22. Little, R. G.; Ibers, J. A. J. Am. Chem. Soc. 1973, 95, 8583.
- 23. Pauling, L. "The Chemical Bond", Cornell University Press: Ithaca, New York, 1967; pp 135-153.
- Buchler, J. W.; Lay, K. Lam; Smith, P. D.; Scheidt, W. R.; Rupperecht,
 G. A.; Kenney, J. E. J. Organomet. Chem. 1976, 110, 109.
- 25. Kirner, J. F.; Dow, W.; Scheidt, W. R. <u>Inorg. Chem.</u> 1976, 15, 1685.
- 26. Cheng, P. T.; Loescher, B. R.; Nyburg, S. C. Inorg. Chem. 1971, 10, 1275.
- 27. Stynes, H. C.; Ibers, J. A. Inorg. Chem. 1971, 10, 2304.
- 28. Fergusson, J. E.; Love, J. L.; Robinson, W. T. Inorg. Chem. 1972, 11, 1662.
- 29. Bright, D.; Ibers, J. A. Inorg. Chem. 1969, 8, 1078.
- 30. Cullen, D. L.; Meyer, E. F., Jr. Acta Crystallogr. Sect. B. 1973, 29, 2507.
- 31. Collins, D. M.; Countryman, R.; Hoard, J. L. <u>J. Am. Chem. Soc.</u> 1972, 94, 2066.
- Hopf, F. R.; O'Brien, T. P.; Scheidt, W. R.; Whitten, D. G. J. Am. Chem. Soc., 1975, 97, 277.
- 33. Kobayashi, T.; Kurokawa, F.; Ashida, T.; Uyeda, N.; Suito, E. <u>J. Chem. Soc.</u>, <u>Chem. Comm.</u> 1971, 1963.
- 34. Freidel, M. K.; Hoskins, B. F.; Martin, R. L.; Mason, S. A. <u>J. Chem. Soc.</u>, Chem. Comm. 1970, 400.
- Kobayashi, T.; Ashida, T.; Uyeda, N.; Suito, E.; Kakudo, M. <u>Bull Chem.</u>
 <u>Soc. Japan</u> 1971, 44, 2095.
- Fischer, M. S.; Templeton, D. H.; Zalkin, A.; Calvin, M. J. Am. Chem. Soc. 1971, 93, 2622.
- 37. Ukei, K. Acta Crystallogr., Sect. B. 1973, 29, 2290.
- 38. Cullen, D. L.; Meyer, E. F., Jr.; Smith, K. M. Inorg. Chem. 1977, 16, 1179.
- 39. Scheidt, W. R.; Dow, W. J. Am. Chem. Soc. 1977, 99, 1101.
- 40. Rogers, D.; Osborn, R. S. J. Chem. Soc., Chem. Comm. 1971, 840.
- 41. Hrung, C. P.; Tsutsuí, M.; Cullen, D. L.; Meyer, E. F., Jr.; Morimoto, C. N. J. Am. Chem. Soc. 1978, 100, 6068.

TABLE I

Crystal Data for $(\dot{c}_{38}^{H}_{21}^{N}_{8})os^{a}$

a = 11.966(9) A	Fw = 809.9
b = 15.705(3) A	z = 4
c = 17.749(5) A	d calcd = 1.721
	$\mu = 43.9 \text{ cm}^{-1} (MoK \alpha \text{radiation})$
$B = 107.92 (4)^{\circ}$	Systematic absences $h0l(h+l)$ odd) $0k0(k)$ odd)
	Space group P 2 ₁ /n
$V = 3125(4) A^3$	

^aIn this and subsequent tables the estimated standard deviation of the least significant figures shown in parentheses.

Table II Fractional Coordinates and Thermal Motion Parameters Derived from the Least-Squares Refinement

Title C

U(23) 61(6)	-	31(111)		=	15(8)			=	-8(10)	-	11(13)	21(13)	4(15)	1(11)	2(11)	-2(11)	-1(11)	-	17(15)	7(14)	-2(12)	-9(11)	15(10)	-5(11)	-	-	2(1	J	=	=	-	= :	(11)2	٠.	: :	-	-2(11)	111161	(6)1-	-14(12)	3(14)	-14(13)	-24(12)	8(12)	-33(13)	26 (01)65-	•
m			21(7)	22(7)		33(8)	25(8)	2(1	37(10)	1 18	86(10)	51(12)	40(11)	-	19	-	100	8(1	-	0	25(10)	19(11)	(01)9	29(9)	21	_	5	0	2(1	211	-	8	19(10)		. ~	-	29(10)	14(10)	23(8)	14(10)	14(14)	0(13)	58(12)	49(10)	(6)99	26(10)	
-	=	= :	-5(10)	=	(6)9-	13(9)	5	=	7:1	-	-	-20(15)	-	36.1	-14(12)	-4(12)	10(13)	33(14)	61(14)	-	16(15)	10(15)	111111	-9(11)	=	-	0(13)	-	-	2	130	-	8(12)			5	-6(111)	11(12)	-15(10)	-15(12)	-6(14)	-14(14)	-13(14)	-19(13)	-80(14)	-35(11)	
276(2)	-		23(4)		28(4)	43(5)	24(4)		. (5)92	46(6)	•			S	_	34(6)		39(7)	65(8)	(8)99	35(6)	36(6)	_	29(5)	_	_	_	_	_	_	0		400	736		_	44(6)	29(6)	27(5)	40(6)	(8)	48(7)	41(6)	55(7)	27(5)	67(5)	
N			47(6)		_	28(5)	47(6)	38(7)	27(6)	_	_	_	39(7)	34(7)	30(7)	30(7)	34(7)	411 73	26(8)	52(8)	49(8)	25(6)	23(6)	36(7)	22(6)	38(7)	_				26(6)	32(7)	29(7)		416 81	_	33(7)	31(6)	37(5)	37(7)	46(8)	32(7)	36(7)	32(6)	65) 26	(9) 169	
U(11) 352(2)	_		32(5)		38(5)	33(5)	37(5)	48(7)	22 (1)	_	81(8)	(6)96	62(8)	20(1)	20 20	40(1)	51(7)	62(8)	46(7)	(2)15	34(6)	46(7)	38(6)	28(6)	37(6)	47(6)	16(7)	11(8)	40(7)	58(7)				34(0)	38(7)	48(7)	29(6)	43(7)	37(5)	31(6)	52(8)	61(8)	84(9)	44(6)	20(6)	62 (1)	
0.11752(3)	.04546	0.0286	0.0896(4)	1914(21436	0.2739(5)	0.1521(4)	0.0084(6)	-0.0625(6)	-0.1073(6)	-0-1704(6)	-0.1889(6)	.1463(-0.0826(5)	-0.0237(6)	. 0.0282(6)	0.0268(6)	-0.0210(6)	-0.0091(7)	0.0492(7)	0.0985(6)	0.0874(6)	0.1269(5)	0.2311(6)	0.3057(6)	0.3493(6)	0.4189(6)	4	0.4014(6)	.3308(.27076	-2180	.22091	0 25637 23	• -		0.1533(6)	-	0.1988(5)	0.2054(6)	0.2613(7)	0.3133(7)	0.3062(6)	0.2449(6)	0.05096 61	0.0109(5)	
0.11171(3)			0.2557(5)			-0.0112(5)	0.0620(6)	0.16311 73	0.21516 71					0.2568(7)	0.2290(7)	0.2300(7)	0.2601(7)	0.3177(7)	0.3290(8)	0.2830(8)	0.2267(7)	0.2150(7)	0.1622(7)	0.06691 7)	0.0206(6)	0.0057(7)	-0.0424(7)	-0.07521 71	-0.0603(7)			0.01050 71		10 000000	-0.04526 73	0.0044(7)	0.0193(7)	6.000011 71	0.2178(5)	0.2481(7)	0.3067(8)	0.3387(7)	0.3112(7)	0.2520(7)	0.0225(8)	-0.0374(5)	
X 0.22473(4)			0.0855(7)	_		0.3804(7)	0.3844(7)	0.3667(9)	0.3569(10)	0.4347(10)	0.3991(111	0.2923(11)	-	0.2483(9)	0.1908(9)	0.03000 91	-0.0855(10)	-0.1664(10)	-0.2715(10)	-0.3023(10)	-0.22441 91	-0.1160(110)	-0.0135(9)	0.08931 91	0.1005(9)	0.02261 91	0.0643(10)	0.1763(111	0.2542(9)	0.2153(10)	-		0.5508(9)	16 19550	-	0.0756(10)	0.5704(9)	0.46601 91	0.3026(7)	0.4043(9)	0.4642(111)	0.4107(11)	0.2990(111)	0.2474(10)	0.1629(9)	0-1273(8)	
ALTA	NC13	N(V)	2 2 2	N(O)N	Niol	N(7)	N(B)	(1)	(7))	((3)	(4)	(5)	(9))	223	(19)	(6)2	((1))	(111)	(71))	((1))	(\$11)	((1))	(011)	(211)	(6113)	((1))	(07)7	(17)3	(77))	(17)	(67) 7	((7))	(07))	(17)	(97)	((10)	(151)	(175)	(6)N	(133)	() () ()	((1))	(0())	C(1)	((1))	000	

^aThe Debye-Waller Factor is defined as $T=\exp[-2\pi^2(U_{11}a*^2h^2+U_{22}b*^2k^2+U_{33}c*^21^2+U_{12}a*b*hk+U_{13}a*c*hl+U_{33}b*c*kl)]$. The values for U have been multiplied by 10^3 , except for those of Os, which have been multiplied by 10^4 .

Table VI. Comparison of CO Frequencies of MP(CO)(Py)

PcM(CO)(Py)	RuPc	RuTPP	OsPc	OsTPP	Os0EP
vC =0 (cm ⁻¹)	1965	1939	1930	1920	1902

P = Pc, TPP, and OEP

Pc = phthalocyanine

TPP = tetraphenylporphine

OEP = octaethylporphyrin

Py = pyridine

M = Os or Ru

TABLE VII $^{\circ} \text{Bond Lengths (A) and Angles (deg)}^{\text{ a}}$

Os-N(2)	2.027(9)		N(2) - Os - N(4)	90.1(4)
Os-N(4)	1.983(9)	2.01(3)	N(2) - Os - N(8)	89.6(4)
0s-N(6)	2.034(9)		N(4) - Os - N(6)	90.1(4) 89.9(3)
0s-N(8)	1.978(9)		N(6) - Os - N(8)	89.7(4)
Os-N(9)	2.202(9)		N(2) - Os - N(6)	174.2(4) 174.3(2)
0s-C(38)	1.83(1)		N(4) - Os - N(8)	174.5(4)
N(2)-C(1)	1.38(1)		N(2) - Os - N(9)	88.2(4)
N(2)-C(8)	1.36(1)		N(4) - Os - N(9)	88.8(4)
N(4)-C(9)	1.39(1)		N(6) - Os - N(9)	86.0(4) 87(1)
N(4)-C(16)	1.37(1)	1.37(1)	N(8)-Os-N(9)	85.7(4)
N(6)-C(17)	1.37(1)	_,,,,,,,	N(2) - Os - C(38)	91.9(5)
N(6)-C(24)	1.36(1)		N(4) - Os - C(38)	92.3(5)
N(8)-C(25)	1.38(1)		N(6) - Os - C(38)	93.8(5) 93(1)
N(8)-C(32)	1.39(1)		N(8)-Os-C(38)	93.2(5)
N(1)-C(1)	1.33(1)		N(9)-Os-C(38)	178.9(6)
N(1)-C(32)	1.33(1)		Os-N(2)-C(1)	124.3(8)
N(3)-C(8)	1.35(1)		Os-N(2)-C(8)	124.3(8)
N(3)-C(9)	1.36(1)		0s-N(4)-C(9)	125.8(8) 125(2)
N(5)-C(16)	1.32(1)	1.34(2)	0s-N(4)-C(16)	126.9(7)
N(5)-C(10)	1.32(1)		Os-N(6)-C(17)	123.3(8)
N(3) - C(24)	1.35(1)		Os-N(6)-C(24)	123.8(9)
N(7)-C(24) N(5)-C(25)	1.37(1)		Os-N(8)-C(24) Os-N(8)-C(25)	126.4(8)
C(1)-C(2)	1.46(1)		Os-N(8)-C(32)	127.3(8)
C(7)-C(8)	1.47(1)		C(1)-N(2)-C(32)	111(1)
			C(1)-N(2)-C(8) C(9)-N(4)-C(16)	
C(9)-C(10)	1.45(2)	1 (6(2)		
C(15)-C(16)	1.47(1)	1.46(2)	C(17)-N(6)-C(24)	113(1)
C(17)-C(18)	1.46(1)		C(25)-N(8)-C(32)	106(1)
C(23)-C(24)	1.47(2)		a(1) (1/1) a(1)	
C(25)-C(26)	1.48(2)	•	C(1)-N(1)-C(32)	125(1)
C(31)-C(32)	1.42(2)		C(8)-N(3)-C(9)	126(1) 125(1)
C(2)-C(7)	1.40(2)		C(16)-N(5)-C(17)	124(1)
C(10)-C(15)	1.42(1)	1.41(1)	C(24)-N(7)-C(25)	125(1)
C(18)-C(23)	1.41(2)		N(1)-C(1)-N(2)	128(1)
C(26)-C(31)	1.40(2)		N(2)-C(8)-N(3)	125(1)
C(2)-C(3)	1.40(2)		N(3)-C(9)-N(4)	126(1)
C(6)-C(7)	1.38(1)		N(4)-C(16)-N(5)	125(1) 127(2)
C(10)-C(11)	1.40(2)		N(5)-C(17)-N(6)	128(1)
C(14)-C(15)	1.38(2)	1.39(1)	N(6)-C(24)-N(7)	129(1)
C(18)-C(19)	1.39(1)		N(7)-C(25)-N(8)	127(1)
C(22)-C(23)	1.39(1)		N(8)-C(32)-N(1)	125(1)
C(26)-C(27)	1.37(2)		N(1)-C(1)-C(2)	125(1)
C(30)-C(31)	1.39(2)		N(3)-C(8)-C(7)	123(1)
C(3)-C(4)	1.38(2)		N(3)-C(9)-C(10)	124(1)
C(5)-C(6)	1.37(2)		N(5)-C(16)-C(15)	124(1)
C(1)-C(12)	1.35(2)		N(5)-C(17)-C(18)	125(1) 124(1)
C(13)-C(14)	1.38(2)	1.38(1)	N(7)-C(24)-C(23)	124(1)
C(19)-C(20)	1.39(2)	1.30(1)	N(7)-C(25)-C(26)	123(1)
C(21)-C(22)	1.36(2)		N(1)-C(32)-C(31)	124(1)
C(27)-C(128)	1.38(2)			
C(29)-C(130)	1.39(2)			

		N(2)-C(1)-C(2)	107(1)
		N(2)-C(8)-C(7)	108(1)
C(4)-C(5)	1.40(2)	N(4)-C(9)-C(10)	110(1)
C(12)-C(13)	1.39(2) 1.39(1)	N(4)-C(16)-C(15)	111(1)
C(20)-C(21)	1.40(2)	N(6)-C(17)-C(18)	106(1) 109(2)
C(28)-C(29)	1.39(2)	N(6)-C(24)-C(23)	106(1)
N(9)-C(33)	1 32(1)	N(8)-C(24)-C(25) N(8)-C(25)-C(26)	110(1)
N(9) - C(33) N(9) - C(37)		N(8)-C(23)-C(26) N(8)-C(32)-C(31)	
	1.36(2)	C(1)-C(2)-C(3)	111(1)/
C(33)-C(34)	1.36(2)		108(1)
C(34)-C(35)	1.36(2) 1.36(1)	C(2)-C(7)-C(8)	106(1)
C(35)-C(36)	1.37(2)	C(9)-C(10)-C(15)	107(1)
C(36)-C(37)	1.37(2)	C(10)-C(15)-C(16)	105(1)
C(38)-0(1)	1.17(1)	C(17)-C(18)-C(23)	107(1) 107(1)
N(2)-N(4)	2.84(1)	C(18)-C(23)-C(24)	107(1)
N(2)-N(8)	2.82(1) $2.83(1)$	C(25)-C(26)-C(31)	105(1)
N(4)-N(6)	2.04(1)	C(26)-C(31)-C(32)	107(1)
N(6)-N(8)	2.83(1)	C(1)-C(2)-C(3)	131(1)
N(2)-N(6)	4.06(1) 4.01(7)	C(6)-C(7)-C(8)	132(1)
N(4)-N(8)	3.90(1)	C(9)-C(10)-C(11)	134(1)
N(1)-N(5)	6.78(1) $6.78(1)$	C(14)-C(15)-C(16)	134(1) 132(1)
N(3)-N(7)	6.77(1)	C(17)-C(18)-C(19)	132(1)
		C(22)-C(23)-C(24)	132(1)
		C(25)-C(26)-C(27)	132(1)
		C(30)-C(31)-C(32)	133(1)
		C(7)-C(2)-C(3)	121(1)
		C(2)-C(7)-C(6)	121(1)
		C(15)-C(10)-C(11)	119(1) 121(1)
		C(10)-C(15)-C(14)	121(1)
		C(23)-C(18)-C(19)	121(1)
		C(18)-C(23)-C(22)	120(1)
		C(31)-C(26)-C(27)	122(1)
		C(26)-C(31)-C(30)	120(1)
		C(2)-C(3)-C(4)	116(1)
		C(5)-C(6)-C(7)	118(1)
		C(10)-C(11)-C(12)	120(1) 118(1)
		C(13)-C(14)-C(15)	119(1)
		C(18)-C(19)-C(20)	117(1)
		C(21)-C(22)-C(23)	119(1)
		C(26)-C(27)-C(28)	118(1)
		C(29)-C(30)-C(31)	118(1)
		C(3)-C(4)-C(5)	122(1)
		C(4)-C(5)-C(6)	121(1)
		C(11)-C(12)-C(13)	121(1)
		C(12)-C(13)-C(14)	121(1) 121(1)
		C(19)-C(20)-C(21)	122(1)
		C(20)-C(21)-C(22)	121(1)
		C(27)-C(28)-C(29)	121(1)
		C(28)-C(29)-C(30)	121(1)
		Os-N(9)-C(33)	122.2(9) 121.8(5)
		Os-N(9)-C(37)	121.5(8)
		C(33)-N(9)-C(38)	116(1)
		N(9)-C(33)-C(34)	124(1) 123(1)
		N(9)-C(37)-C(36)	123(1)
		C(33)-C(34)-C(35)	120(1)
		C(34)-C(35)-C(36)	118(1) 119(1)
		C(35)-C(36)-C(37)	119(1)
		0s-C(38)-O(1)	177(1)

a. Some nonbonded distances of interest are also given. Figures in parenthesis for the averaged values are the root-mean-square standard deviations of the least significant figure

TABLE VIII
Least-Squares Planes

A. Deviations (a) from planes

Plane 1 Plane 2 Plane 3 Plane 4 Plane 5 Plane 6 Plane 7								
N(1)		Plane 1					Plane 6	Plane 7
N(2) -0.006	0s	-0.154	-0.099		-0.056	0.017	-0.070	
N(3)	N(1)	-0.082					0.088	
N(4) -0.075 -0.003 -0.090 -0.018 0.251 0.039 N(5) -0.157 -0.029 -0.143 0.102 0.102 -0.158 N(6) -0.003 0.014 0.180 -0.013 -0.013 -0.117 N(7) 0.067 0.156 0.288 0.424 -0.053 -0.009 N(8) -0.041 -0.004 0.113 0.096 -0.028 0.011 C(1) -0.049 -0.068 0.014 -0.176 0.148 0.160 C(2) -0.029 -0.075 -0.011 -0.287 0.261 0.258 C(3) -0.042 -0.121 -0.016 -0.390 0.240 0.295 C(4) 0.030 -0.069 0.000 -0.437 0.418 0.437 C(5) 0.117 0.030 0.028 -0.379 0.615 0.546 C(6) 0.097 0.043 0.001 -0.309 0.603 0.478 C(7) 0.020 -0.013 -0.022 -0.267 0.420 0.329 C(8) 0.014 0.016 -0.018 -0.157 0.389 0.258 C(9) 0.025 0.078 -0.047 -0.034 0.462 0.209 C(10) 0.070 0.144 -0.054 0.018 0.600 0.255 C(11) 0.159 0.228 -0.031 0.021 0.813 0.397 C(12) 0.111 0.205 -0.115 0.009 0.830 0.333 C(13) -0.051 0.075 -0.250 -0.031 0.612 0.102 C(14) -0.118 0.013 -0.253 -0.011 0.422 -0.021 C(15) -0.062 0.044 -0.159 0.010 0.412 0.052 C(16) -0.102 0.001 -0.130 0.035 0.244 -0.031 C(17) -0.124 0.002 -0.048 0.196 0.018 -0.166 C(18) -0.037 0.118 0.086 0.416 0.014 -0.157 C(19) -0.066 0.122 0.044 -0.159 0.010 0.412 0.052 C(20) 0.039 0.246 0.205 0.244 0.031 0.012 C(21) 0.157 0.353 0.386 0.845 -0.004 -0.106 C(22) 0.187 0.351 0.423 0.787 0.021 -0.026 C(22) 0.083 0.128 0.046 0.295 0.041 0.042 -0.023 C(21) 0.157 0.353 0.386 0.845 -0.004 -0.106 C(22) 0.083 0.122 0.411 0.412 -0.223 0.011 C(24) 0.021 0.128 0.195 0.387 -0.016 -0.055 C(25) 0.005 0.060 0.216 0.256 -0.093 -0.012 C(27) 0.083 0.122 0.411 0.412 -0.220 0.012 C(27) 0.083 0.122 0.411 0.412 -0.220 0.001 C(29) -0.010 -0.027 0.328 0.159 -0.329 0.008 C(20) -0.099 -0.094 0.070 -0.041 -0.106 -0.003 C(33) -0.079 -0.094 0.070 -0.041 -0.106 -0.003 C(33) -0.079 -0.094 0.070 -0.041 -0.106 -0.003 C(33) -0.099 -0.094 0.070 -0.041 -0.106 -0.003 C(33) -0.0099 -0.094 0.070 -0.041 -0.106 -0.003	N(2)	-0.006						
N(5) -0.157 -0.029 -0.143 0.102 0.102 -0.158 N(6) -0.003 0.014 0.180 -0.013 -0.013 -0.017 N(7) 0.067 0.156 0.288 0.424 -0.053 -0.009 N(8) -0.041 -0.004 0.113 0.096 -0.028 0.011 C(1) -0.049 -0.068 0.014 -0.176 0.148 0.160 C(2) -0.029 -0.075 -0.011 -0.287 0.261 0.258 C(3) -0.042 -0.121 -0.016 -0.390 0.240 0.295 C(4) 0.030 -0.069 0.000 -0.437 0.418 0.437 C(5) 0.117 0.030 0.028 -0.379 0.615 0.546 C(6) 0.097 0.043 0.001 -0.309 0.603 0.478 C(7) 0.020 -0.013 -0.022 -0.267 0.420 0.329 C(8) 0.014 0.016 -0.018 -0.157 0.389 0.258 C(9) 0.025 0.078 -0.047 -0.034 0.462 0.209 C(10) 0.070 0.144 -0.054 0.018 0.600 0.255 C(11) 0.159 0.228 -0.031 0.021 0.813 0.397 C(12) 0.111 0.205 -0.115 0.009 0.830 0.333 C(13) -0.051 0.075 -0.250 -0.031 0.612 0.102 C(14) -0.118 0.013 -0.253 -0.011 0.422 -0.021 C(15) -0.062 0.044 -0.159 0.010 0.412 0.052 C(16) -0.102 0.001 -0.130 0.035 0.244 -0.031 C(17) -0.124 0.002 -0.048 0.196 0.018 -0.166 C(18) -0.037 0.118 0.086 0.416 0.014 -0.157 C(19) -0.066 0.122 0.001 -0.130 0.035 0.244 -0.031 C(17) -0.124 0.002 -0.048 0.196 0.018 -0.166 C(18) -0.037 0.118 0.086 0.416 0.014 -0.157 C(19) -0.066 0.122 0.001 -0.130 0.035 0.244 -0.031 C(27) 0.083 0.226 0.205 0.244 0.542 0.001 -0.236 C(20) 0.039 0.246 0.208 0.699 -0.012 -0.236 C(21) 0.157 0.353 0.386 0.845 -0.004 -0.106 C(22) 0.187 0.351 0.423 0.787 0.0021 -0.236 C(22) 0.187 0.351 0.423 0.787 0.0021 -0.236 C(22) 0.187 0.351 0.423 0.787 0.0021 -0.006 C(24) 0.021 0.128 0.195 0.387 -0.016 -0.055 C(25) 0.005 0.060 0.216 0.256 -0.093 0.008 C(22) 0.083 0.122 0.411 0.317 -0.333 0.008 C(22) 0.009 -0.004 0.070 -0.041 -0.106 -0.003 N(9)	N(3)	0.053	0.074		-0.111		0.297	
N(6) -0.003 0.014 0.180 -0.013 -0.013 -0.117 N(7) 0.067 0.156 0.288 0.424 -0.028 0.011 C(1) -0.049 -0.068 0.014 -0.176 0.148 0.160 C(2) -0.029 -0.075 -0.011 -0.287 0.261 0.258 C(3) -0.042 -0.129 -0.075 -0.011 -0.390 0.240 0.295 C(4) 0.030 -0.069 0.000 -0.437 0.418 0.437 C(5) 0.117 0.030 0.028 -0.379 0.615 0.546 C(6) 0.097 0.043 0.001 -0.399 0.603 0.478 C(7) 0.020 -0.013 -0.022 -0.267 0.420 0.329 C(8) 0.014 0.016 -0.018 -0.157 0.389 0.258 C(9) 0.255 0.078 -0.047 -0.034 0.462 0.209 C(9	N(4)	-0.075	-0.003	-0.090	-0.018	0.251	0.039	
N(7)	N(5)	-0.157	-0.029	-0.143	0.102	0.102	-0.158	
N(8)	N(6)	-0.003	0.014	0.180	-0.013	-0.013	-0.117	
C(1)	N(7)	0.067	0.156	0.288	0.424			
C(2) -0.029 -0.075 -0.011 -0.287 0.261 0.258 C(3) -0.042 -0.121 -0.016 -0.390 0.240 0.295 C(4) 0.030 -0.069 0.000 -0.437 0.418 0.437 C(5) 0.117 0.030 0.028 -0.379 0.615 0.546 C(6) 0.097 0.043 0.001 -0.309 0.603 0.478 C(7) 0.020 -0.013 -0.022 -0.267 0.420 0.329 C(8) 0.014 0.016 -0.018 -0.157 0.389 0.258 C(9) 0.025 0.078 -0.047 -0.034 0.462 0.209 C(10) 0.070 0.144 -0.054 0.018 0.600 0.255 C(11) 0.159 0.228 -0.031 0.021 0.813 0.397 C(12) 0.111 0.205 -0.115 0.009 0.830 0.333 C(13) -0.051	N(8)	-0.041	-0.004	0.113	0.096	-0.028	0.011	
C(3)	C(1)	-0.049	-0.068	0.014	-0.176	0.148	0.160	
C(4)	C(2)	-0.029	-0.075	-0.011	-0.287	0.261	0.258	
C(4)	C(3)	-0.042	-0.121	-0.016	-0.390	0.240	0.295	
C(5) 0.117 0.030 0.028 -0.379 0.615 0.546 C(6) 0.097 0.043 0.001 -0.309 0.603 0.478 C(7) 0.020 -0.013 -0.022 -0.267 0.420 0.329 C(8) 0.014 0.016 -0.018 -0.157 0.389 0.258 C(9) 0.025 0.078 -0.047 -0.034 0.462 0.209 C(10) 0.070 0.144 -0.054 0.018 0.600 0.255 C(11) 0.159 0.228 -0.031 0.021 0.813 0.397 C(12) 0.111 0.205 -0.115 0.009 0.830 0.333 C(13) -0.051 0.075 -0.250 -0.031 0.612 0.102 C(14) -0.118 0.013 -0.253 -0.011 0.422 -0.021 C(15) -0.062 0.044 -0.159 0.010 0.412 0.052 C(16) -0.102 0.001 -0.130 0.035 0.244 -0.031 C(17) -0.124 0.002 -0.048 0.196 0.018 -0.166 C(18) -0.037 0.118 0.086 0.416 0.014 -0.157 C(19) -0.066 0.122 0.049 0.478 -0.009 -0.236 C(20) 0.039 0.246 0.208 0.699 -0.012 -0.203 C(21) 0.157 0.353 0.386 0.845 -0.004 -0.106 C(22) 0.187 0.351 0.423 0.787 0.021 -0.026 C(23) 0.062 0.205 0.244 0.542 0.001 -0.080 C(24) 0.021 0.128 0.195 0.387 -0.016 -0.055 C(25) 0.005 0.060 0.216 0.256 -0.093 -0.012 C(26) 0.014 0.049 0.279 0.259 -0.178 -0.003 C(27) 0.083 0.122 0.411 0.412 -0.232 0.011 C(28) 0.046 0.059 0.411 0.437 -0.333 -0.008 C(30) -0.079 -0.102 0.194 0.003 -0.275 -0.007 C(31) -0.052 -0.049 0.184 0.069 -0.183 0.003 C(33) -0.079 -0.102 0.194 0.003 -0.275 -0.007 C(31) -0.052 -0.049 0.184 0.069 -0.183 0.003 C(33) -0.009 -0.094 0.070 -0.041 -0.106 -0.003 C(33) -0.0099 -0.094 0.070 -0.041 -0.106 -0.003 C(33) -0.0099 -0.094 0.070 -0.041 -0.106 -0.003 C(33) -0.0099 -0.094 0.070 -0.0041 -0.106 -0.003 C(33) -0.0099 -0.094 0.070 -0.0041 -0.106 -0.003		0.030	-0.069	0.000	-0.437	0.418	0.437	
C(6) 0.097 0.043 0.001 -0.309 0.603 0.478 C(7) 0.020 -0.013 -0.022 -0.267 0.420 0.329 C(8) 0.014 0.016 -0.018 -0.157 0.389 0.258 C(9) 0.025 0.078 -0.047 -0.034 0.462 0.209 C(10) 0.070 0.144 -0.054 0.018 0.600 0.255 C(11) 0.159 0.228 -0.031 0.021 0.813 0.397 C(12) 0.111 0.205 -0.115 0.009 0.830 0.333 C(13) -0.051 0.075 -0.250 -0.031 0.612 0.102 C(14) -0.118 0.013 -0.253 -0.011 0.422 -0.021 C(15) -0.062 0.044 -0.159 0.010 0.412 0.052 C(16) -0.102 0.001 -0.130 0.035 0.244 -0.031 C(17) -0.124 0.002 -0.048 0.196 0.018 -0.166 C	C(5)	0.117	0.030	0.028	-0.379	0.615	0.546	
C(7)		0.097			-0.309	0.603	0.478	
C(8)				-0.022	-0.267	0.420		
C(9) 0.025 0.078 -0.047 -0.034 0.462 0.209 C(10) 0.070 0.144 -0.054 0.018 0.600 0.255 C(11) 0.159 0.228 -0.031 0.021 0.813 0.397 C(12) 0.111 0.205 -0.115 0.009 0.830 0.333 C(13) -0.051 0.075 -0.250 -0.031 0.612 0.102 C(14) -0.118 0.013 -0.253 -0.011 0.422 -0.021 C(15) -0.062 0.044 -0.159 0.010 0.412 0.052 C(16) -0.102 0.001 -0.130 0.035 0.244 -0.031 C(17) -0.124 0.002 -0.048 0.196 0.018 -0.166 C(18) -0.037 0.118 0.086 0.416 0.014 -0.157 C(19) -0.066 0.122 0.049 0.478 -0.009 -0.236 C(20) 0.039 0.246 0.208 0.699 -0.012 -0.203 C(21) 0.157 0.353 0.386 0.845 -0.004 -0.106 C(22) 0.187 0.351 0.423 0.787 0.021 -0.026 C(23) 0.062 0.205 0.244 0.542 0.001 -0.080 C(24) 0.021 0.128 0.195 0.387 -0.016 -0.055 C(25) 0.005 0.060 0.216 0.256 -0.093 -0.012 C(26) 0.083 0.122 0.411 0.412 -0.232 0.011 C(28) 0.046 0.059 0.411 0.337 -0.333 -0.008 C(29) -0.010 -0.027 0.328 0.159 -0.329 0.008 C(30) -0.079 -0.102 0.194 0.003 -0.275 -0.007 C(31) -0.052 -0.049 0.184 0.069 -0.183 0.003 C(32) -0.099 -0.094 0.070 -0.041 -0.106 -0.003 N(9)					-0.157	0.389		
C(10) 0.070 0.144 -0.054 0.018 0.600 0.255 C(11) 0.159 0.228 -0.031 0.021 0.813 0.397 C(12) 0.111 0.205 -0.115 0.009 0.830 0.333 C(13) -0.051 0.075 -0.250 -0.031 0.612 0.102 C(14) -0.118 0.013 -0.253 -0.011 0.422 -0.021 C(15) -0.062 0.044 -0.159 0.010 0.412 0.052 C(16) -0.102 0.001 -0.130 0.035 0.244 -0.031 C(17) -0.124 0.002 -0.048 0.196 0.018 -0.166 C(18) -0.037 0.118 0.866 0.416 0.014 -0.157 C(19) -0.066 0.122 0.049 0.478 -0.009 -0.236 C(20) 0.039 0.246 0.208 0.699 -0.012 -0.203 C(21) 0.157 0.353 0.386 0.845 -0.004 -0.106 C(22) 0.187 0.351 0.423 0.787 0.021 -0.026 C(23) 0.062 0.205 0.244 0.542 0.001 -0.080 C(24) 0.021 0.128 0.195 0.387 -0.016 -0.055 C(25) 0.005 0.060 0.216 0.256 -0.093 -0.012 C(26) 0.014 0.049 0.279 0.259 -0.178 -0.003 C(27) 0.083 0.122 0.411 0.412 -0.232 0.011 C(28) 0.046 0.059 0.411 0.337 -0.333 -0.008 C(29) -0.010 -0.027 0.328 0.159 -0.329 0.008 C(30) -0.079 -0.102 0.194 0.003 -0.275 -0.007 C(31) -0.052 -0.049 0.184 0.069 -0.183 0.003 C(32) -0.099 -0.094 0.070 -0.041 -0.106 -0.003 N(9)								
C(11) 0.159 0.228 -0.031 0.021 0.813 0.397 C(12) 0.111 0.205 -0.115 0.009 0.830 0.333 C(13) -0.051 0.075 -0.250 -0.031 0.612 0.102 C(14) -0.118 0.013 -0.253 -0.011 0.422 -0.021 C(15) -0.062 0.044 -0.159 0.010 0.412 0.052 C(16) -0.102 0.001 -0.130 0.035 0.244 -0.031 C(17) -0.124 0.002 -0.048 0.196 0.018 -0.166 C(18) -0.037 0.118 0.086 0.416 0.014 -0.157 C(19) -0.066 0.122 0.049 0.478 -0.009 -0.236 C(20) 0.039 0.246 0.208 0.699 -0.012 -0.203 C(21) 0.157 0.353 0.386 0.845 -0.004 -0.106 C(22) 0.187 0.351 0.423 0.787 0.021 -0.026 C(23) 0.062 0.205 0.244 0.542 0.001 -0.080 C(24) 0.021 0.128 0.195 0.387 -0.016 -0.055 C(25) 0.005 0.660 0.216 0.256 -0.093 -0.012 C(26) 0.014 0.049 0.279 0.259 -0.178 -0.003 C(27) 0.083 0.122 0.411 0.412 -0.232 0.011 C(28) 0.046 0.059 0.411 0.337 -0.333 -0.008 C(29) -0.010 -0.027 0.328 0.159 -0.329 0.008 C(30) -0.079 -0.102 0.194 0.009 -0.275 -0.007 C(31) -0.052 -0.049 0.184 0.069 -0.183 0.003 C(32) -0.099 -0.094 0.070 -0.041 -0.106 -0.003 N(9)		0.070						
C(12) 0.111 0.205 -0.115 0.009 0.830 0.333 C(13) -0.051 0.075 -0.250 -0.031 0.612 0.102 C(14) -0.118 0.013 -0.253 -0.011 0.422 -0.021 C(15) -0.062 0.044 -0.159 0.010 0.412 0.052 C(16) -0.102 0.001 -0.130 0.035 0.244 -0.031 C(17) -0.124 0.002 -0.048 0.196 0.018 -0.166 C(18) -0.037 0.118 0.086 0.416 0.014 -0.157 C(19) -0.066 0.122 0.049 0.478 -0.009 -0.236 C(20) 0.039 0.246 0.208 0.699 -0.012 -0.203 C(21) 0.157 0.353 0.386 0.845 -0.004 -0.106 C(22) 0.187 0.351 0.423 0.787 0.021 -0.026 C(23) 0.062 0.205 0.244 0.542 0.001 -0.080 C(24) 0.021 0.128 0.195 0.387 -0.016 -0.055 C(25) 0.005 0.060 0.216 0.256 -0.093 -0.012 C(26) 0.014 0.049 0.279 0.259 -0.178 -0.003 C(27) 0.083 0.122 0.411 0.412 -0.232 0.011 C(28) 0.046 0.059 0.411 0.337 -0.333 -0.008 C(29) -0.010 -0.027 0.328 0.159 -0.329 0.008 C(30) -0.079 -0.102 0.194 0.003 -0.275 -0.007 C(31) -0.052 -0.049 0.184 0.069 -0.183 0.003 C(33) -0.079 -0.102 0.194 0.003 -0.275 -0.007 C(31) -0.052 -0.049 0.184 0.069 -0.183 0.003 C(33) -0.099 -0.094 0.070 -0.041 -0.106 -0.003 N(9)			0.228	-0.031				
C(14) -0.118			0.205	-0.115	0.009	0.830		
C(14) -0.118				-0.250	-0.031	0.612		
C(15) -0.062				-0.253	-0.011	0.422	-0.021	
C(16) -0.102			0.044	-0.159	0.010			
C(17) -0.124	C(16)							
C(18) -0.037								
C(19) -0.066					0.416			
C(20) 0.039 0.246 0.208 0.699 -0.012 -0.203 C(21) 0.157 0.353 0.386 0.845 -0.004 -0.106 C(22) 0.187 0.351 0.423 0.787 0.021 -0.026 C(23) 0.062 0.205 0.244 0.542 0.001 -0.080 C(24) 0.021 0.128 0.195 0.387 -0.016 -0.055 C(25) 0.005 0.060 0.216 0.256 -0.093 -0.012 C(26) 0.014 0.049 0.279 0.259 -0.178 -0.003 C(27) 0.083 0.122 0.411 0.412 -0.232 0.011 C(28) 0.046 0.059 0.411 0.337 -0.333 -0.008 C(29) -0.010 -0.027 0.328 0.159 -0.329 0.008 C(30) -0.079 -0.102 0.194 0.003 -0.275 -0.007 C(31) -0.052 -0.049 0.184 0.069 -0.183 0.003 C(32) -0.099 -0.094 0.070 -0.041 -0.106 -0.003 N(9)								
C(21) 0.157 0.353 0.386 0.845 -0.004 -0.106 C(22) 0.187 0.351 0.423 0.787 0.021 -0.026 C(23) 0.062 0.205 0.244 0.542 0.001 -0.080 C(24) 0.021 0.128 0.195 0.387 -0.016 -0.055 C(25) 0.005 0.060 0.216 0.256 -0.093 -0.012 C(26) 0.014 0.049 0.279 0.259 -0.178 -0.003 C(27) 0.083 0.122 0.411 0.412 -0.232 0.011 C(28) 0.046 0.059 0.411 0.337 -0.333 -0.008 C(29) -0.010 -0.027 0.328 0.159 -0.329 0.008 C(30) -0.079 -0.102 0.194 0.003 -0.275 -0.007 C(31) -0.052 -0.049 0.184 0.069 -0.183 0.003 C(32) -0.099 -0.094 0.070 -0.041 -0.106 -0.003 N(9)								
C(22)								
C(23)	C(22)	0.187						
C(24) 0.021 0.128 0.195 0.387 -0.016 -0.055 C(25) 0.005 0.060 0.216 0.256 -0.093 -0.012 C(26) 0.014 0.049 0.279 0.259 -0.178 -0.003 C(27) 0.083 0.122 0.411 0.412 -0.232 0.011 C(28) 0.046 0.059 0.411 0.337 -0.333 -0.008 C(29) -0.010 -0.027 0.328 0.159 -0.329 0.008 C(30) -0.079 -0.102 0.194 0.003 -0.275 -0.007 C(31) -0.052 -0.049 0.184 0.069 -0.183 0.003 C(32) -0.099 -0.094 0.070 -0.041 -0.106 -0.003 N(9)	C(23)	0.062	0.205	0.244	0.542	0.001	-0.080	
C(25) 0.005 0.060 0.216 0.256 -0.093 -0.012 C(26) 0.014 0.049 0.279 0.259 -0.178 -0.003 C(27) 0.083 0.122 0.411 0.412 -0.232 0.011 C(28) 0.046 0.059 0.411 0.337 -0.333 -0.008 C(29) -0.010 -0.027 0.328 0.159 -0.329 0.008 C(30) -0.079 -0.102 0.194 0.003 -0.275 -0.007 C(31) -0.052 -0.049 0.184 0.069 -0.183 0.003 C(32) -0.099 -0.094 0.070 -0.041 -0.106 -0.003 N(9)								
C(26) 0.014 0.049 0.279 0.259 -0.178 -0.003 C(27) 0.083 0.122 0.411 0.412 -0.232 0.011 C(28) 0.046 0.059 0.411 0.337 -0.333 -0.008 C(29) -0.010 -0.027 0.328 0.159 -0.329 0.008 C(30) -0.079 -0.102 0.194 0.003 -0.275 -0.007 C(31) -0.052 -0.049 0.184 0.069 -0.183 0.003 C(32) -0.099 -0.094 0.070 -0.041 -0.106 -0.003 N(9)	C(25)	0.005				-0.093	-0.012	
C(27) 0.083 0.122 0.411 0.412 -0.232 0.011 C(28) 0.046 0.059 0.411 0.337 -0.333 -0.008 C(29) -0.010 -0.027 0.328 0.159 -0.329 0.008 C(30) -0.079 -0.102 0.194 0.003 -0.275 -0.007 C(31) -0.052 -0.049 0.184 0.069 -0.183 0.003 C(32) -0.099 -0.094 0.070 -0.041 -0.106 -0.003 N(9)			0.049	0.279	0.259	-0.178		
C(28)	C(27)		0.122	0.411	0.412	-0.232		
C(29) -0.010 -0.027 0.328 0.159 -0.329 0.008 C(30) -0.079 -0.102 0.194 0.003 -0.275 -0.007 C(31) -0.052 -0.049 0.184 0.069 -0.183 0.003 C(32) -0.099 -0.094 0.070 -0.041 -0.106 -0.003 N(9)					0.337	-0.333	-0.008	
C(30) -0.079 -0.102 0.194 0.003 -0.275 -0.007 C(31) -0.052 -0.049 0.184 0.069 -0.183 0.003 C(32) -0.099 -0.094 0.070 -0.041 -0.106 -0.003 N(9)								
C(31) -0.052 -0.049 0.184 0.069 -0.183 0.003 C(32) -0.099 -0.094 0.070 -0.041 -0.106 -0.003 N(9)	C(30)	-0.079	-0.102	0.194	0.003			
C(32) -0.099 -0.094 0.070 -0.041 -0.106 -0.003 N(9)			-0.049				0.003	
N(9) 0.020 C(33) 0.013 C(34) 0.004 C(35) 0.005								
C(33)		*						0.020
C(34) -0.004 C(35) 0.013 C(36) -0.005								
C(35) = = 0.013 C(36) - 0.005								
C(36)0.005		= ==						
		·• · · · · ·		-				
	C(37)							-0.011

B. Angles (deg) between Least-Squares Planes

	Plane 2	Plane 3	Plane 4	Plane 5	Plane 6	Plane 7
Plane 1	1.4	2.7	5.2	5.1	3.1	98.6
Plane 2		3.0	4.1	5.5	4.2	98.0
Plane 3			4.1	7.9	5.3	96.1
Plane 4				9.4	8.3	94.1
Plane 5					3.7	103.4
Plane 6						101.4

C. Equations of Planes

Plane 1 Marcocycle;
$$N(1)N(8)$$
, $C(1)-C(32)$
2.721x + 12.314y + 8.382z = 3.126

Plane 2 Isoindole nitrogen atoms;
$$N(2)$$
, $N(4)$, $N(6)$, $N(8)$ 2.509x + 12.216y + 8.726z = 3.053

^aAll planes are unweighted. x,y,z are in monoclinic fractional coordinates.

C(16)-O(1) ^I	3.08	N(3)-C(35) II	3.32
0(1)-0(1) ^I	3.18	N(5)-C(4) III	3.40
$N(4)-O(1)^{I}$	3.25	C(23)-C(35) IV	3.41
$C(15)-O(1)^{I}$	3.26	C(24)-C(36) IV	3.42
$C(38)-O(1)^{I}$	3.31		

^aRoman numeral superscripts denote the following equivalent positions relative to the reference molecule at x,y,z:

I - x, - y, - z
II -
$$\frac{1}{2}$$
 + x, $\frac{1}{2}$ - y, - $\frac{1}{2}$ + z
III - $\frac{1}{2}$ + x, $\frac{1}{2}$ - y, $\frac{1}{2}$ - z
IV 1 - x, - y, - z





Fig. 1 Tsutsui

Figure 1. Coordination of carbon monoxide by PcRu(II) in THF. The peak at 627 nm is characteristic for PcRu(II) and the peaks at 640 and 580 nm are due to PcRu(CO)(THF). Upon coordination of carbon monoxide, the peak at 627 disappears, while new peaks (at 540 and 580 nm) appear.

Tsutsuf



is shown. The bond lengths involving the osmium ion are also shown. The thermal ellipsoids are drawn for 50% probability, except those of ORTEP¹³ drawing of the structure of PcOg(CO)(Py). Numbering scheme Figure 2.




Figure 3. Stereoview of PcOs(II)(CO)(Py)




Fig. 4

Figure 4. Stereoview of the packing in the unit cell. The osmium atoms are indicated by the larger circles, the nitrogen atom. Oxygen atoms of the carbonyl groups are labeled with the letter 0.

SYNTHESIS AND STRUCTURE OF A NEW CLASS

OF METALLOPHTHALOCYANINES:

Carbonylphthalocyanato(pyridine or THF)ruthenium(II) and carbonylphthalocyanato
(pyridine or THF)osmium(II)

Shuichi Omiya and Minoru Tsutsui Department of Chemistry Texas A&M University College Station, Texas 77843

Edgar F. Meyer, Jr.

Department of Biophysics and Biochemistry
Texas A&M University
College Station, Texas 77843

Ivan Bernal
Department of Chemistry
University of Houston
Houston, Texas 77004

David L. Cullen
Department of Chemistry
Connecticut College
New London, Connecticut 06320

TABLE III

Calculated Postional Parameters for Hydrogen Atoms

Atom	x	<u>¥</u>	<u>z</u>
H(3)	0.512	0.197	-0.094
H(4)	0.444	0.292	-0.205
H(5)	0.274	0.368	-0.232
H(6)	0.142	0.342	-0.159
H(11)	-0.145	0.351	-0.064
H(12)	-0.324	0.369	-0.040
H(13)	-0.378	0.292	0.055
H(14)	-0.244	0.198	0.140
H(19)	-0.054	0.028	0.333
H(20)	0.015	-0.053	0.450
H(21)	0.203	-0.108	0.492
H(22)	0.331	-0.083	0.419
H(27)	0.618	-0.090	0.319
H(28)	0.796	-0.117	0.293
H(29)	0.832	-0.055	0.181
H(30)	0.691	0.028	0.093
H(33)	0.449	0.227	0.168
H(34)	0.544	0.327	0.264
H(35)	0.451	0.379	0.356
H(36)	0.256	0.334	0.342
H(37)	0.167	0.233	0.244

ROOT-HEAN-SQUARE AMPLITUDES OF THEFMAL VIBRATION IN AMSSTROMS.

ATOM	min.	INT'MED.	MAX.
57777777777777777777777777777777777777	979407357167665648114989902588867718510922198924750	77820915250535919622120065765767651341546586251146 79797979747987062000000077779657676513415465465867071116 1111111111111111111111111111111	03979798-78-9040950-90005-15407001-8866886084848889 901-390144459859000570751-15407094800505050508608 1000000000000000000000000000000000

Values of 10 | Fo | and 10 | Fc |
For PcOs(CO)(Py)

PAGE C	etere damad
#	
90E	
\mathbf{C}	
6000 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	673 673 1159 8151 815
######################################	25.558 35.558
r eesseeeeeeeeeeeeeee	
7-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1	
# # # # # # # # # # # # # # # # # # #	%888 <u>49</u>
T normodiano-prodictar-perode-panaro-dianapor	- m = m r
5	- 21212121
######################################	22222
0	14223 1723 1733 1733 1733 1733 1733 1733 1
608	52854
- 5 - Fe-4480Fe5-7777776-6480Fe5-77777-6-78480Fe6-5-7	エいーロエ
$-\frac{\pi}{2}$. While is a state of the state of the substitution of	CONNE
	50000
7. P. C.	17 4 9 1 9 2 8 4 7 8
K	134845
	20079
	MMMMM.
#	002203

ずいたのだいこうりゅんんかよししっしいないのとのここにらっかんりっしょとっつこうかんしょ 8000 **8888499778778778778788888488** COPPOSITOR OF CONTRACTOR OF CONTRACTOR OF CONTRACTOR OF THE PROPERTY OF CONTRACTOR OF

	FCALC	1016 392	972	4:	2023	479	2433	20.	16.4.3	151	165	205.3	170	50.70	77.4	100 100 100 100 100 100 100 100 100 100	637	391	716 619	1628	803	1767	183	936	955	200	1086	200	16.00	1.1.1	£.	9 ja	5 12	213
	FOBS	995 368	277	125	2076	151	2356		7000	505	300	2019	153	7020	750	900	653	134	713 638	16-40	837	1817	20.7	9.41	196	205	1037	1.57	12.90	27	10.33	0 - .	13 13 13 13	215
	ر د د	=======================================	2-12	1		2 - 3	CIC	٥.	-0	110	70	n w		200	120	1	2-12	3-12	= 7	9	2 6	71	77	7	0.	- 0	110	٠ د د د	2.0	r-	00	==	27	9 15
M	ī	22	nn nn	101	้าเก	01	N		10	101	CTC	un	101	10	101	n	va	2															ou co	C1
PAGE																																		
P.	FCALC	326	1031	1065	1653	1550	986	1870	1373	499	836	1780	427	2111	1571	362	371	812	504 846	423	1210	1252	1028	403	1316	1163	1723	1072	200	187.4	10-11	1459	1462 684	405
	FOBS	361	1023	1105	1675	1606	666	1934	1351	517	839	17.45	400	979	1514	275	39.1	822	525	7	181	1259	10.58	777	1060	1219	1752	1082	1 13	1761	1020		19.7 12.7 13.7	105E
	ن	<u>v. 9</u>	1-15	=:	2 0	2	9	1,	d id	7	70	-	(4)	·) <	רטיז	ဖ	J. C	2	95	m	<u> </u>	9	2 1	٩	ن ا	7 (-7	-	o -	• 6	m	7	םי ה	~ €	Ξ
	~	44	ທ່າທ່	ம்	ກ່ທ	רט	in i	s	រាម) u)	ט כט		, ,	2. 6	, ,	u ı ı	.) u	1 113	m ıt	Ψ	ω α	·w	ש ע	σ.	ω.	ם ים	Œ	w (د، د	פי	121	در و	1012	פי
	I	an	NV	ומו	หณ	יח	CU	N	NO	114	NI	u c	101	NC	111	N	v.c.	10	NN	101	M.C.	ini	NC	ın	NI	40	161	es e	. c.	CI	C1 C	M C:	61.01	
	FCALC	358 1161	2251	294	12:10	238	310	369	25025	1350	993	37.1	734	252	834	109	1000 1000 1000 1000 1000 1000 1000 100	530	600	37.4	1916 255	2:107	272	512	1838	2, 5, 1, 5, 5, 5, 5, 5, 5, 5, 5, 5, 5, 5, 5, 5,	1922	761	; ;	1321		- C3	4 15 4 15	7.45
	FOBS	1217	2321	315	1269	241	218	332	2401	1274	915	100	712	206	263	623	24.0 7.0 7.0 7.0 7.0	529	611	359	37.6	2.156	1229	513	1855	9.5	1037	1773	7.7	0.:21		500	5.50 5.70 5.70	652
F.	٦	47	יש ער !	41	24	-	9	-,	.1 1.5	7	ភេទ	٠,	. ෆ	ש פ	27	2:	2	13	27	2	22	တု၊	~ Y	יש	71	20	0		. • •	7	Ų.	۰ <u>ء</u>	===	Ξ
11 7	¥	M (6)	m r	ו ניא (دما زم	19	L -2 (M) I	W W	110	M) (0.10	(11	w) (w) t•J	60.1	~ h	4	7 4	+	+ +	4	1 5	4	Τ,	1 4	=	÷ •	-	-;	-	: 1		7
FILE	Ξ		Nr	141	7 C!	N	N	C) (101	C) C	u (101	20	iu	N:	'u r.	C.	n n	(0)	NN	(1)	N C	in	C) (V.V	101	C1 C	. * .	CI	C4:	۰, ۸۱	C. M	24
114 FO	FCHLC	1595	6. 5. 6. 5. 6. 5. 6. 5. 7. 5.	26.43	7.2	10.58	16.31	192	1097	10.17	664	15.5	203	1,20	1 4	623	1439	610	1492 1361	1338	338	201	307	1032	2073	15.5	323	(19		=	63.1	200	991-	1100
HUMELL	FOBS	1652	263	1984	710	1560	107.4	204	10.00	1042	673	19	632	188	1 1 1 1	619	5.53	603	761-1	26-13	2033	520	1.035	363	13:1	15.27	311			7:15	02.2		2017 2011	=======================================
300	۷	ដូដូ	7 =	- 0	14	S	اد	~ (: c	. 01	215	2 =	9	- 5	2 7	2	3=	9	m #	8 _ 1	e cr	71			l., 0	n r-		Ξ:	: ::	95	-:	I	==	P
FIRE	¥					-	-			-			-	1,0	ı çı	'n	116	i,	יוט	(1)	M (1	010	, r	101	2	u N	C:	٠, ٢		.h	ė.		10 to	160
23	Ξ	~ (1	NN	101	101	CI	~:	-11	vu	101	U) C	u n	N	40	1 61	L1 (VN	101	un	1.01	U C	CI (NC	101	n: n	A CI	CI	010		. 71	er:	161	OLC:	
(PYR LD ME) (CREBBILD) PHTRLGC/RIB 100SHTURGTE)	FCHLC	411	767	5	6.15 1.00	200	213		7 65	611	328	306	1001	203	367	507	925	391	2400	505	3601	1677	557	1301	10 to 00 to	100 100 100 100 100 100 100 100 100 100	2008	23.5		1.n.1	610	11.11	1781 1781	2.5
CHURD	FORS	372	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	7	663	730	530		2007	533	00.5	836	933	2 2 2	335	1153	1238	380	2437	803	1167	1631	5071	1225	33.1	7	6115	7] I.	10.23	3		1750	1031
E																																		
=	د	-1	7 1	r- 1		~	çı.		- 11	17	7 0	7	0	V 7	20	9	TO	2	φ 7	CI	3 C	4	0 00	9	<u></u>	10	-	2	: =	17	-	3.7	1.10	7
RID'.	л Г	-m 11	7 K	-	200	15 -3	9	2		15 3	7 tr	7 3	16 0	2 12	0-13	0-16	0-17	0-10	φ 7 Θ =	0	20	0	9 6	9 10	0 0	1-13	1-17	91-1		1-15		= T	' ' '	1 -3

	S FCALC		
	- FOBS		
7	×	wwwwwwqqqqqqqqqqqqqqqqqqqqqqqqqqqqqqqq	
PAGE	U	M T M A A OLT M R UN A M A M A M A M A M A M A M A M A M A	7 N. # (1 M. M.)
•	FCAL		
	FOBS		
	A L	:พุทธเกตเดนตอนตอนตอนตอนตอนตอนตอนตอนตอนตอนตอนตอนตอน	
	I	**************************************	
	FCAL(
	FOBS		
10 FC	A L		-00160606
FO AND 18	T U		
01 (FCAL		
SITUME	FOBS	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	
VR LD (HE) COMPRONYL) PHTALOCYANATOOSNIUMCI I	H K	<u> </u>	1111000
HTALOC	u	A. N. C. M. K P. D. D. D. M. M. M. M. N. C. D. D. D. M.	
UNTLIP	FCAL		
E) (CHPE	FOBS		
RICTOR	7	######################################	
6	I	าเกษาการเกษาการเกษาการเกษาการเกษาการเกษาการเกษาการเกษา	reier ereier ereier

	FCALC	2882-1-128-1
	FOBS	7.7.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2
	л Г	
3 39	_	
. PAGE	FCHLC	80000000000000000000000000000000000000
	FCBS	20000000000000000000000000000000000000
	H 7.	พถดนพมนพมนพมนพมนพมนพมนพมนพมนพมนพมนพมนมนนนนนน
	FCALC	866 866 866 866 866 866 866 866 866 866
	FOBS	80000000000000000000000000000000000000
AND 10 FC	H K	
10 60	FERE	888 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
iture 11:	sac:	2000 2000 2000 2000 2000 2000 2000 200
TUOST	7	
СУЯНЕ	z z	$ \begin{array}{c} WEADMORDER ADMORDER $
EYR IL INES (CHREONYL) PHTHLOCYANGTOOSNIUM (11)	FCALC	25 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
CHEBO	FOES	######################################
FYRILINE	π ۲	

									,																			Olick			m	Mes	Ties	the s	1000	-	-	
	ALC	213	200	536	689	181	100	130	1-13	389	700	000	366	21.5	-	121	0	35.	200	RY I	950	= =====================================	620	515	33	598	170	950	000	50	100	7	; ; ; .	7	3 "		33	13
	FCAL							-							•	-	• "						-			• •			-	-		_		-	_		7.6	
	FOBS	13	800	7	90	9:0	IM	200	157	283	200	100	393	7	9	10	100	7	90	7 12	200	60	919	2 CT	60.	-6	0 m	20	m	200	3	3	-	# :	2 42	==	- 7	9
	3	e	• •	, ,	-	911	•	· =		1.10	.0 .	ניי ט	.,	-					- 1	ני כנ		1 11)	w e	1, 1		ט כט	, 1.,	עיי	. ت		01.	_		-	-	123		
	_	2	= 0	18	2-	3	7 7	!	•	C) I	71	.=	6.	2	-1	-	- ני	e e	ų, ·	- (10	ci	71	١٥٥	9	Ģ	-	N	7.	- 1-	- :-	200	=	7.1		1	- :	
	¥		70	101	2	2:2	72	101	2	M.	710	i W	13	12	2:	3 5	100	Ξ	3	Ξ.	1 1	ব	I :	107	12	<u> </u>	ים ב	12	יירט		9 6		-	= 3	5 5	0	= c	2
21	I	4	7	7	77	4.	1 7	: 7	7	4.	7 -	1 7	-	7	7 -	: 7	1	7	₹.	т-	7 7	7	7.	1 7	7	4 -	1 7	ব	-1 -	T U	רט ר	מנ	U~	UT 13	7 67	S)	ט רש	רט ר
PAGE																																						
٩	FCALC	730	561	970	607	545	455	633	503	684	145	47.4	4E3	163	565	200	275	647	733	216	102	024	672	27.4	512	643	510	537	210	200	100	560	405	367	19	3-1-	4 4 4	7 =
	FC					_	-			_	-				•		•				-	_							-		-	•	_					
	FOBS	751	1000	932	624	250	200	120	189	531	580	300	483	975	200	220	332	962	776	313	1 3	927	168	0 7	175	960	0 0	976	70	000	0.00	0.35	503	5	20.7	33.5	100	72
	Œ					=:	•	=		Ξ.	= `										_	.=	-			-			=		-	•	÷	•	•			
	_	10	= 4	30	=			ו ו			n	-	<u>m</u>	C!:	=	7 0) ;	9	, נה		; -	. 2	OI I	n 7	e.	~ 0	0 0	=	<u> </u>	30	0 14	न्	64	= (V 7	Ċ)	د. و	===
	×	œ	တင်	Ġ	ģ	o	ממ	. 0	0	0.0	9	ν ξ	9	3	2	20	0	0	9	2	22	10	0	2 5	2	9	2 2	9	-:	_ :	==	: ::	=	=:	==	=	==	: 2
	Ξ	4	4	1	4	4	4 -	1 7	7	ব	4 .	: 4	7	7	7	7 5	7 7	7	7	7 -	7 7	7	ব :	ד ד	7	₹.	: 1	4	7.	1 -	1 7	7	7	ব •	1	7	ব -	
	FCAL C	893	226	686	8.49	423	0 to 0.	52.1	564	370	139	100	000	253	000	200	30	035	350	150	298	629	480	567	575	627	300	395	102	2000		163	832	000		10	500	100
	F		-						-	-	•		_	_		-		_								•	-	-	-		-	-						
	FOBS	873	203	633	833	427	900	200	132	393	899	177	7	363	600	200	ייים	916	364	1 C	27.2	628	475	2017	575	525	200	135	113	25	12	133	316	7			215	2 2
	IF		-				•	•	-	-	•	-	=	-	-			=								•	-	_	-		-	-						
FC	٦	ဖ	r- a	9	=	12	1 :	1 =	2	9	ب ا	7 1	Ç.	0	٠1،	4 R	υ	ò	D i	9 .	2 19	2	2	<u>''-</u>	10	60	נו ב	i,	41	20		- c	O)	M.	7 .7		r- :	: :-
10 6	¥	()	w u	a	ω	(0)	ם ר	- : -	-1-	۱-۱	r- r	- 1-	٠.	N-1		-1	- 1 -	٠.	<u>~</u> 1		- 2	φ.	က် (က် ထဲ	တ်	00 0	(C)	ယ	ω (o c	0 00	m	ಾ	0.0	0 0	32	C. r	C
=	z	4	7-	7	4	4	¢ <	1 7		4	7.	1 1	7	ব	÷ :	4 ~	7	4	1		1 5	77	4.	र ग	4	- ,	र प	7	4.	1.	- 7	7	-	•	: =	•		• -
FO H						_																																
10 F	נטרכ	1133	7 9	0	912	9.0	000	187	153	339	6.1	7 00 0 00 0 0	100	562	3	20.00		27.0	643	-		- 9	633	5 2 2 2	100		200	15	9	110		3	1:	Ξ	-			
2	L								_								•		_															•				
3	FOBS	138	Či z	8253	861	10 1		38.5	110	3.42	652	710	163	568	900	40 P	137	181	5.5	200	106	1-	31.0	2007	583	70	100	160	182		200	1.	00	=		100	1 - 1	
UIL	Ŀ	-							-		•	-	-		N			1	-												6	1		•	VI.		•	•
[00]	_	9	=:	: 7	-	5		20	=	01.	5	- 16	רטי	7	m :	7 -		. VI	S	w 0	0 5	10	=!	0 H	12	13	20	=	011	0 1	- 47	7	•••	Çi •	- ::	-	::	-
E E	¥	प	7 ,	7	רע	ו רע	n d	לש ל	'n	n,	ע כע	רע מ	165	57	so i	n v	רט כ	רטי	ر ا	0	ט גר	0	9	n is	ú	δ.	6	פֿיי	10 1	با د	o u	υ	e	i i	0 :	10	12 1	: 2
E .	I	4	4 4	4	T	7	7 -	7	~	4	4 .	1 7	7	4	7 .	7 7	1 1	4	ন :	7 -	1 4	7	4.	1 7	7	7 .	: 4	4	7.	-	7 7	-	7	• •	7	7		: -
1																																						
Ē	FCALC	312	333	153	102	522		28.5	15.	999	300	100	3	130	0.6	000	128	33	383	22.	610	133	562		-	348	369	E71	SUS:	1	1	100	200	J :	-			
J.O.F.	FC							-			-		-			-		_	-			_		-	-	,	,	-			-		64	•	•	-	-	
F.F.	FCRS	31.	9 2	20:	320	9	0.00	0 m	7	5. i	0.0	199	130	**, : 	37.0	91	5	1.	21	7 1	1 00	513	600	T 12	199	77	21.0	933	6.	1	11:	1	53			25		
PARTOTHES CLEREAINESPHARECYRNATOOSHIUMCLD	Ĭ.							-			-		-		= :	-	-	-			•	-	- :	-	-		1	=			-		3.	•		~	-	
1:IE	١	33	0 5	?=	<u></u>	97 5	2 1	- 7	2	= :	2 0	n :-		7	., .		. 2	-	٠,٠	7 8	w	00	0	111	91	2	3.6	00		0 -	7 1	-	=	- 1		~	w:	- 33
E.	2	C1	rı r	114	r,	NI		. 4.	4	-	5	רא ני	10	100	0	. ,.	m	~	iu i	w 14	רא ר	*	ו רא	0 ;0	+	+ -	+ +	7	4.		7 17	7	•		: -:	••	ψ: :	7
3	I	7	7 7	7	7	•••	7 -	7 7	7	7	7 .	1 7	7	7	7 .	7 7	7	7	7	7 -	-	7	7.	1 7	7	ч-	17	•	٠,٠	7 -	7 -:	7	-	7 .	• ••	7		•

FCALC	524486686644646675646 864686884686468666866866	1054
FOBS	11000000000000000000000000000000000000	1000
- × • ±	$\omega_0\omega_0\omega_0\omega_0\omega_0\omega_0\omega_0\omega_0\omega_0\omega_0\omega_0\omega_0\omega_0\omega$	_
PAGE		385
FOBS	22000000000000000000000000000000000000	755
Ξ Ά	$ \frac{1}{2} 1$	ี เก
FCALC	4404476668886886866668456858888888888888888	100
FOE5	488414282888888888888888888888888888888	603
10 FC K	<u> </u>	5
FO BNU 10	មស្រាស្សស្រាស្សស្រាស្សស្រាស្សស្រាស្សស្រាសស្រាសស្រាសស្រាស្សស្រាស្សស្រាស្សស្រាស្សស្រាស្សស្រាស្សស្រាស្សស្រាស្សស្	
ECE!		1.04
FORS	25.22.22.22.22.22.22.22.22.22.22.22.22.2	1.5
ILINATUOS K L	$ \begin{array}{c} \mathbf{M} \mathbf{M} \mathbf{M} \mathbf{M} \mathbf{M} \mathbf{M} \mathbf{M} M$	7
THEOUGH H	សស្សាស្ត្រស្ត្រស្ត្រស្ត្រស្ត្រស្ត្រស្ត្រ	w ⁻
INT. FR	600	11:3
FARES	Bundar garage and an	-
PEREDIMENTAL AND LENGTHANDE AND RETURNATION IN THE FORS	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	7 11

ID INE	CARPU	(CARPOAYL) PHTALOCYRIGETOOSMIUN (T)	1.007	三	10	ารเกาน	£110	10 FO H	HIND II	7							PAGE	න න				
	S	FCALC	I	¥			F085 .	CALC	I	~	r Fn	roes F	FCALC	=	ж -	FORS	FCALC	I	~	د	FOBS	FCH
	1691	1967	ωų		M -		336	202	0 1	10 10		733	712		•	727	706	ויםי	03 0	84	893 893	88
	336	503	ω	-			202	1217	oω	מי	8	03	1078	. · ·	2-12	744	797	w	ω	'n	1170	=
	632	612	٤	-	v		:03	390	ور	m		36	462		7	533	546	9	8	4	830	8
	0100	391	w u		0		700	197	יםי	 		430	426		7	543	545	6 ()	တ၀	200	967	43
	31.	254	שנ	• -			205	- C	0 10	7		9 9	196		1	0000	534	ם יב	οα	1-	213	60
	7	141	9 (6	-	=		000	021) W	-		13	503		-	563	558	פינ	000	.0	801	83
	130	503	0	-	-		663	500	פיי	1-1		36	1-103		٠,	1499	14-18	9	8	N	610	7-3
	433	460	9	CI	-1		3.7	1.12	w	4-1			633		1	838	834	9	8	m	754	75
	7 7	113	ω	2	1		159	366	9	1		17	695		1	360	349	۵۰	ဆ	IJ	413	4
	C1.	837	9	CI	-		101	1106	פי	1		31	1501		٠	9+1	431	Þ	တ	و	1011	; J.
	47.5	1617	9	V			505	COL	Ü	1		52	12:42			1577	1555	(L)	သ	۲-	613	13
	808	893	e	CI	1		356	1863	عا	1		90	489		1	620	603	2	3	10	783	733
	1001	990	9	2	-		392	293	ω	1		65	1172		5- 5	585	612	۵.	9	-13	924	93
	633	631	9	CI	1		52	1754	ω	1		63	546		1- 5	1085	1126	ניי	Ġ	12	552	ST
	653	703	9	CI	i		150	218	w	4	2 10	-13	9201			603	585	٩	Ġ	=-	1172	116
15 -2	1:57	417	9	CI	1		655	664	ω	1	1	- 15	61.5			723	729	(L)	σ.	6-	F1-7	57
	2111	1134	9	C.1	1		520	17.29	و،	7	0	102	467			1395	1429	9	9	۲-	10-11	160
	1.1.	77.0	w	C			17.7	193	2	4	1 4	104	395			562	410	9	a)	10	1452	177
	1:15	585	9	CI	111		27.3	2.02.1	2	4	31 5	10	1645			531	533	9	6	-	346	100 C 1
	5.5	576	9	N	• '		13	259	o	4	3	77	529			675	704	9	σ.	-	1173	121
	2115	2159	9	U			163	12:2	0'	7		121	856		2 9	226	976	9	6	'n	365	65
	1.75	9.13	9	r:			593	556	9	7		4	1260			427	443	t,	o,	S	1.109	142
	3 ac.	953	Ü	L1	-		. 673	552	ı,	7		1-01	381	9		909	596	ם	on :	-1	520	อรู
	070	31-81	9	(4)	-		-1-1	197	9	= :		12	737		-	452	217	(O)		σ.	026	5.
	2006	0.25.1	O.	107	-		5.5	503	د،	5		90	6.42		_	223	999	ים		1	555	3
	1166	1163	ا ت	•••	-		113	727	œ	-		æ ;	1193		7	777	1504	0		-10	0001	56
	3	2	υ,	•••	-		303	3	2	-	示i す!	10	533	ים	21-1	of S	5655	ים		2	010	7 0
	3	125.0	ינ	• 1	-		20 1	165	۰	-		20	916		ı	1597	1586	ים	2	7. C	200	5 7
	1353	1265	0	1 (*)	-		5.2	500	י פי	- i		23	16.18			6/2	9-19	۱ ۵	2	D 1	901	
	3 :	200	D (•1	-		333	010	2 (-		٥.	500		1		21.0	2	3	ם נ	0 - 5 0 2 0	200
	0	91	: ن		-		200	G (1)	ات	וה		ا د	700		1	0000	10-10	. د ه	20	0	1000	
	7	90	٥.	0.1	. 1		9:	מממ	۰ م	î Du		0:	101	٥	1	12:2	250	0 1	2 9	1	200	7.7
	100	7.7.	0	0 1			200	00	ى د				101	01	91	000	000	0 1	2 5		2 2 2	
	000	- 0	0 (• •	7		3:	757	٥.	ī D L	٥.	000	9.7	0 (10	0.00	100	י ס.	2 9	- (
		5.	۵	*1	1		113	7.00	_	(7	70	(-)	۰	5	201	15-71	י פ	2	41	100	200
	31,1	6.10.	ů.		1		22	901	2	(1)	3 15	20	1505	ع	-	336	410	2	=	•	65.	-
	21.3		E.	1	1		30.	56.1	. 5	l G	VI M	15	100		CJ	មា () ()	860	2	3	9	7.1-	C)
	12.5	1927	3		1		7	51511	ء.	5	iri D	9	368		च :-	33.0	215	۵۰	=		- E	
	**		120		-		:35	1.11	<i>i</i> :	L'I	0 -	1.3	13.45		ω 1-	7.76	976	12	Ξ	c: -	~:	==
		5.15	2	***			1	1175	12	١-		-	Stre		8-15	725	612	3	Ξ	= -	1.5	15.
	15.00	77.7	ت				-:	53533		Ľ.	3	1:1	:101:	د،		= -	436	10	=	ç	2,12	
	11:5:1	11:1	10	210				5.751	9	.71	7	35	21.1	'n	3-13	07.0	608	9	=	Œ.	1130	I-r.u
	1.67.	17.7	.T.	**)			253	.65	.0		01 5	96.1	11	10	01-3	1056	1021	وا	=	7!	111.7.2	101
		1113		10			==	5: 1-1	12	u	-1	10	.185	כי	8-16	M 12 5	514	42	=	c:	10.1.1	107.
	150%	15.11	10				1.1	0.3	12	-	. 8	7.4	614	.2	1	758	755		=	=	7.	15.
																					5	
)	

e PAGE 9	FCALC	
	FOBS	20000000000000000000000000000000000000
	H K L	
	FCALC	28831 28821
	FOBS	• • • • • • • • • • • • • • • • • • •
	= ×	
	FCALC	25.5 25.5 25.5 25.5 25.5 25.5 25.5 25.5
	FUBS	8.00.00.00.00.00.00.00.00.00.00.00.00.00
IN FO PHO 180 FC	H T. L	 หายหายหายหายหายหายหายหายหายหายหายหายหายห
	FCAL.C	2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2
THE P	FOBS	69974427
CHIMTOUSE	II K	サントトトトトトトトトトトトトトトトトトトトトトトトトトトトトトトトトトトト
(FYREE) THE CHEBOTHE PHINEDCHINGTOUSHINGTOLD	FCALC	
	FOOT	
CFT RID INED	H	$\begin{array}{llllllllllllllllllllllllllllllllllll$

	FCALC	948.74.00
	F085	887-100-100-100-100-100-100-100-100-100-10
	ж г	พอดออออออออออออ
PAGE 10	I	$oldsymbol{a}$ and old
	FCALC	800007488708888888848888888888888888888
	FOBS	- 1888 -
	H K	$ \begin{array}{c} \mathbf{o} \mathbf{o} \mathbf{o} \mathbf{o} \mathbf{o} \mathbf{o} \mathbf{o} o$
	FCALC	12422222222222222222222222222222222222
	FOES	64464 64
19 FO AILE 10 FC	нкг	$ \begin{array}{c} \mathbf{x} \mathbf{u} \mathbf{u} \mathbf{u} \mathbf{u} \mathbf{u} \mathbf{u} \mathbf{u} u$
	FCALC	450 50 50 50 50 50 50 50 50 50 50 50 50 5
HIJIMITE I I	FOBS	######################################
AT00SF	ير بر	เทตเทตเดเตเพลดผลผลผลผลผลผลผลผลผลผล 44444444444444444
DC 7AN	I	ဘလးသက္လက္သက္သက္သက္သက္သက္သက္သက္သက္သည္သည္သည္သည္သည္သည္သည္သည္သည္သည္သည္သည္သည္
TIL) PHTRLE	FCALE	######################################
CHPLA):	FOES	EBS 18 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
(PTR.J.D.HIC) CONFLOCION PHTRLOCIANATOOSHIUMCIT)	H K	

. PAGE 11	FCALC	######################################
	FOBS	100 100 100 100 100 100 100 100 100 100
	KL	************************************
	Έ	
	FCALC	888 883 883 883 883 883 883 883 883 883
	FOBS	8888474588848486888888484888888888888888
	H K L	6000000000000000000000000000000000000
	FCALC	######################################
	FOBS	888445684466844688868888888888888888888
PYRIDINE) (CASECUTY, PHTALOCYANATOOSIJUNCTI) 10 FO HAB 18 FC	ب ب	
	Ξ	••••••••••••••••••••••••••••••••••••
	FCALC	8034555555555555555555555555555555555555
	FURS	28224427862862442426628442662844266284426868844268688844268688888888
	7	คลดลดหรรรรรรษฐติดตอดตอดตอดตอดตอดตอดตอดตอดตอด แพล :มาพพมที่อื่อติดต่อ ไม่เปกพมที่อื่อต้องสอดติดหังสำกับ-กร <i>า</i> กม
	I	ភាគាគល់ សាលាសាលាសាលាសាលាសាលាសាលាសាលាសាលាសាលាសាល
	FCALC	8
	FOES	Range Brand
(PYRIDINE)	1 H	ัง กลง พลง คลง คลง คลง คลง คลง คลง คลง คลง คลง ค

	FCALC	
	FOES	
	_	
	<u>~</u>	
PACE 12	Ξ	
	LCUIC	
	FOBS	
	_	
	4	
	=	
	FCAI C	
	FOBS	
	7	
FC	- ×	
0 10	=	
IN TO AMP TO FC		MS 210 NO - 10 A - 10 A - 2 A A A A A A A A A A A A A A A A A
	FCALC	2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2
Hurre 11	Foos	
TUOS	_	######################################
HILL	I	
HLUC		
MLIPH	FCHLC	
EVETDING COARGONAL) PROBLOGYBRING (1)	FUBS	######################################
DITE.	۰	. ๆกางเหมนุยมแพนพมพมพมมาวุวกระบบระบบตามจากของจากการการการการการ อาการการบับทุกตามปายมหายอังหากับมา-กล่างประจับก็ชับการการกำหน้า-
PYFII		ักการและสารายการและ เลือน การการการการการการการการการการการการการก
3		